

Approximation of variationally posed eigenvalue problems

Daniele Boffi

Dipartimento di Matematica “F. Casorati”

Università di Pavia

`daniele.boffi@unipv.it`

`http://www-dimat.unipv.it/boffi/`

14-16 May 2007

Plan of the lecture

- ☞ Standard finite element approximation of Laplace eigenproblem

Plan of the lecture

- ☞ Standard finite element approximation of Laplace eigenproblem
- ☞ Approximation theory of compact operators

Plan of the lecture

- ☞ Standard finite element approximation of Laplace eigenproblem
- ☞ Approximation theory of compact operators
- ☞ The curl curl eigevalue problem

Plan of the lecture

- ☞ Standard finite element approximation of Laplace eigenproblem
- ☞ Approximation theory of compact operators
- ☞ The curl curl eigenvalue problem
- ☞ Laplace eigenproblem in mixed form

Plan of the lecture

- ☞ Standard finite element approximation of Laplace eigenproblem
- ☞ Approximation theory of compact operators
- ☞ The curl curl eigenvalue problem
- ☞ Laplace eigenproblem in mixed form
- ☞ A counterexample (inf-sup OK and spurious eigenvalues)

Plan of the lecture

- ➡ Standard finite element approximation of Laplace eigenproblem
- ➡ Approximation theory of compact operators
- ➡ The curl curl eigenvalue problem
- ➡ Laplace eigenproblem in mixed form
- ➡ A counterexample (inf-sup OK and spurious eigenvalues)
- ➡ Finite element approximation of eigenproblems in mixed form

Plan of the lecture

- ☞ Standard finite element approximation of Laplace eigenproblem
- ☞ Approximation theory of compact operators
- ☞ The curl curl eigenvalue problem
- ☞ Laplace eigenproblem in mixed form
- ☞ A counterexample (inf-sup OK and spurious eigenvalues)
- ☞ Finite element approximation of eigenproblems in mixed form

Variationally posed eigenproblem (Laplace operator)

Eigenvalue problem

Find $\lambda \in \mathbb{R}$ such that for some $u \in V$ with $u \neq 0$ it holds

$$a(u, v) = \lambda b(u, v) \quad \forall v \in V$$

Variationally posed eigenproblem (Laplace operator)

Eigenvalue problem

Find $\lambda \in \mathbb{R}$ such that for some $u \in V$ with $u \neq 0$ it holds

$$a(u, v) = \lambda b(u, v) \quad \forall v \in V$$

Assumptions

$$H \quad (= L^2(\Omega)) \quad , \quad V \quad (= H_0^1(\Omega)) \quad \subset H$$

Hilbert spaces, V compactly embedded in H

$$a(u, v) \quad (= \int_{\Omega} \nabla u \cdot \nabla v \, d\mathbf{x}) \quad V \times V \rightarrow \mathbb{R}$$

bilinear, continuous, symmetric, coercive

$$b(u, v) \quad (= (u, v)) \quad H \times H \rightarrow \mathbb{R}$$

bilinear, continuous, symmetric

Laplace eigenproblem

Strong form

$$\begin{aligned} -\Delta u &= \lambda u && \text{in } \Omega \\ u &= 0 && \text{on } \partial\Omega \end{aligned}$$

Laplace eigenproblem

Strong form

$$\begin{aligned} -\Delta u &= \lambda u & \text{in } \Omega \\ u &= 0 & \text{on } \partial\Omega \end{aligned}$$

Weak form

$\lambda \in \mathbb{R}$, $u \in V$, $u \neq 0$:

$$a(u, v) = \lambda b(u, v) \quad \forall v \in V$$

Laplace eigenproblem

Strong form

$$\begin{aligned} -\Delta u &= \lambda u && \text{in } \Omega \\ u &= 0 && \text{on } \partial\Omega \end{aligned}$$

Weak form

$\lambda \in \mathbb{R}$, $u \in V$, $u \neq 0$:

$$a(u, v) = \lambda b(u, v) \quad \forall v \in V$$

Resolvent operator

$T : H \rightarrow H$, $T(H) \subset V$ implies T is compact

$$a(Tf, v) = b(f, v) \quad \forall v \in V$$

Laplace eigenproblem

Strong form

$$\begin{aligned} -\Delta u &= \lambda u & \text{in } \Omega \\ u &= 0 & \text{on } \partial\Omega \end{aligned}$$

Weak form

$\lambda \in \mathbb{R}$, $u \in V$, $u \neq 0$:

$$a(u, v) = \lambda b(u, v) \quad \forall v \in V$$

Resolvent operator

$T : H \rightarrow H$, $T(H) \subset V$ implies T is compact

$$a(Tf, v) = b(f, v) \quad \forall v \in V$$

$\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_i \leq \dots$ (numbered s. t. all have multiplicity one)

$E_i = \text{span}(u_i)$, normalization $b(u_i, u_i) = 1$

$V = \bigoplus_{i=1}^{\infty} E_i$ (orthonormal basis)

Laplace eigenproblem: approximation

$$V_h \subset V, \dim V_h = N(h)$$

Discrete problem

Find $\lambda_h \in \mathbb{R}$ such that for some $u_h \in V_h$ with $u_h \neq 0$ it holds

$$a(u_h, v) = \lambda_h b(u_h, v) \quad \forall v \in V_h$$

Laplace eigenproblem: approximation

$$V_h \subset V, \dim V_h = N(h)$$

Discrete problem

Find $\lambda_h \in \mathbb{R}$ such that for some $u_h \in V_h$ with $u_h \not\equiv 0$ it holds

$$a(u_h, v) = \lambda_h b(u_h, v) \quad \forall v \in V_h$$

Discrete (compact) resolvent operator

$$T_h : H \rightarrow H$$

$$a(T_h f, v) = b(f, v) \quad \forall v \in V_h$$

Laplace eigenproblem: approximation

$$V_h \subset V, \dim V_h = N(h)$$

Discrete problem

Find $\lambda_h \in \mathbb{R}$ such that for some $u_h \in V_h$ with $u_h \not\equiv 0$ it holds

$$a(u_h, v) = \lambda_h b(u_h, v) \quad \forall v \in V_h$$

Discrete (compact) resolvent operator

$$T_h : H \rightarrow H$$

$$a(T_h f, v) = b(f, v) \quad \forall v \in V_h$$

$$\lambda_{1,h} \leq \lambda_{2,h} \leq \cdots \leq \lambda_{i,h} \leq \cdots \leq \lambda_{N(h),h}$$

$$E_{i,h} = \text{span}(u_{i,h}), \text{ normalization } b(u_{i,h}, u_{i,h}) = 1$$

$$V_h = \bigoplus_{i=1}^{N(h)} E_{i,h}$$

Rayleigh quotient

Continuous problem

$$\lambda_1 = \min_{v \in V} \frac{a(v, v)}{b(v, v)} = R(v), \quad u_1 = \arg \min_{v \in V} R(v)$$

Rayleigh quotient

Continuous problem

$$\lambda_1 = \min_{v \in V} \frac{a(v, v)}{b(v, v)} = R(v), \quad u_1 = \arg \min_{v \in V} R(v)$$

$$\lambda_i = \min_{v \in \left(\bigoplus_{k=1}^{i-1} E_k \right)^\perp} R(v), \quad u_i = \arg \min_{v \in \left(\bigoplus_{k=1}^{i-1} E_k \right)^\perp} R(v)$$

Rayleigh quotient

Continuous problem

$$\lambda_1 = \min_{v \in V} \frac{a(v, v)}{b(v, v)} = R(v), \quad u_1 = \arg \min_{v \in V} R(v)$$

$$\lambda_i = \min_{v \in \left(\bigoplus_{k=1}^{i-1} E_k \right)^\perp} R(v), \quad u_i = \arg \min_{v \in \left(\bigoplus_{k=1}^{i-1} E_k \right)^\perp} R(v)$$

Discrete problem

$$\lambda_{1,h} = \min_{v \in V_h} R(v), \quad u_{1,h} = \arg \min_{v \in V_h} R(v)$$

$$\lambda_{i,h} = \min_{v \in \left(\bigoplus_{k=1}^{i-1} E_{k,h} \right)^\perp} R(v), \quad u_{i,h} = \arg \min_{v \in \left(\bigoplus_{k=1}^{i-1} E_{k,h} \right)^\perp} R(v)$$

Rayleigh quotient (cont'ed)

A first easy consequence

$$\lambda_1 = \min_{v \in V} R(v), \quad \lambda_{1,h} = \min_{v \in V_h} R(v)$$

$$V_h \subset V \text{ implies } \boxed{\lambda_{1,h} \geq \lambda_1}$$

Rayleigh quotient (cont'ed)

A first easy consequence

$$\lambda_1 = \min_{v \in V} R(v), \quad \lambda_{1,h} = \min_{v \in V_h} R(v)$$

$$V_h \subset V \text{ implies } \boxed{\lambda_{1,h} \geq \lambda_1}$$

Same argument does not apply to the next eigenvalues

$$\lambda_i = \min_{v \in \left(\bigoplus_{k=1}^{i-1} E_k \right)^\perp} R(v), \quad u_i = \arg \min_{v \in \left(\bigoplus_{k=1}^{i-1} E_k \right)^\perp} R(v)$$

$$\lambda_{i,h} = \min_{v \in \left(\bigoplus_{k=1}^{i-1} E_{k,h} \right)^\perp} R(v), \quad u_{i,h} = \arg \min_{v \in \left(\bigoplus_{k=1}^{i-1} E_{k,h} \right)^\perp} R(v)$$

Rayleigh quotient (cont'ed)

An alternative representation

$$\lambda_i = \min_{E \in V_i} \max_{v \in E} R(v)$$

where V_i denotes the set of all subspaces E of V with $\dim E = i$

Rayleigh quotient (cont'ed)

An alternative representation

$$\lambda_i = \min_{E \in V_i} \max_{v \in E} R(v)$$

where V_i denotes the set of all subspaces E of V with $\dim E = i$

Proof

$\min \max \leq \lambda_i$ Take $E = \bigoplus_{k=1}^i E_k$, $v = \sum_{k=1}^i \alpha_k u_k$, use orthogonalities and $\lambda_k \leq \lambda_i$ for all $k \leq i$ to get $R(v) \leq \lambda_i$

Rayleigh quotient (cont'ed)

An alternative representation

$$\lambda_i = \min_{E \in V_i} \max_{v \in E} R(v)$$

where V_i denotes the set of all subspaces E of V with $\dim E = i$

Proof

$$\begin{aligned} \min \max &\leq \lambda_i && \text{Take } E = \bigoplus_{k=1}^i E_k, \ v = \sum_{k=1}^i \alpha_k u_k, \text{ use orthogonalities} \\ &&& \text{and } \lambda_k \leq \lambda_i \text{ for all } k \leq i \text{ to get } R(v) \leq \lambda_i \\ \min \max &\geq \lambda_i && \text{If } E \neq \bigoplus_{k=1}^i E_k \text{ then there exists } v \in E \text{ with } v \perp u_k \\ &&& \text{for all } k \leq i, \text{ hence } R(v) > \lambda_i \end{aligned}$$

Rayleigh quotient (cont'ed)

Continuous min-max

$$\lambda_i = \min_{E \in V_i} \max_{v \in E} R(v)$$

where V_i denotes the set of all subspaces E of V with $\dim E = i$

Discrete min-max

$$\lambda_{i,h} = \min_{E \in V_{i,h}} \max_{v \in E} R(v)$$

where $V_{i,h}$ denotes the set of all subspaces E of V_h with $\dim E = i$

Rayleigh quotient (cont'ed)

Continuous min-max

$$\lambda_i = \min_{E \in V_i} \max_{v \in E} R(v)$$

where V_i denotes the set of all subspaces E of V with $\dim E = i$

Discrete min-max

$$\lambda_{i,h} = \min_{E \in V_{i,h}} \max_{v \in E} R(v)$$

where $V_{i,h}$ denotes the set of all subspaces E of V_h with $\dim E = i$

Consequence

For all $i = 1, \dots, N(h)$ one has $\lambda_{i,h} \geq \lambda_i$

Convergence of eigenvalues/eigenfunctions

Some notation

$m : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\lambda_{m(1)} < \lambda_{m(2)} < \cdots < \lambda_{m(N)} < \cdots$$

Convergence of eigenvalues/eigenfunctions

Some notation

$m : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\lambda_{m(1)} < \lambda_{m(2)} < \cdots < \lambda_{m(N)} < \cdots$$

$\hat{\delta}(E, F) = \max(\delta(E, F), \delta(F, E))$, where E, F subspaces of H

$$\delta(E, F) = \sup_{u \in E, \|u\|_H=1} \inf_{v \in F} \|u - v\|_H$$

Convergence of eigenvalues/eigenfunctions

Some notation

$m : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$\lambda_{m(1)} < \lambda_{m(2)} < \dots < \lambda_{m(N)} < \dots$$

$\hat{\delta}(E, F) = \max(\delta(E, F), \delta(F, E))$, where E, F subspaces of H

$$\delta(E, F) = \sup_{u \in E, \|u\|_H=1} \inf_{v \in F} \|u - v\|_H$$

Definition of convergence

<Comments>

$\forall \varepsilon > 0, \forall N \in \mathbb{N}, \exists h_0 > 0$ such that $\forall h \leq h_0$

$$\blacktriangleright \max_{i=1, \dots, m(N)} |\lambda_i - \lambda_{i,h}| \leq \varepsilon$$

$$\blacktriangleright \hat{\delta} \left(\bigoplus_{i=1}^{m(N)} E_i, \bigoplus_{i=1}^{m(N)} E_{i,h} \right) \leq \varepsilon$$

Convergence of eigenmodes (cont'ed)

Remark: from the monotonicity property, the problem we are studying cannot present spurious eigenvalues

Convergence of eigenmodes (cont'ed)

Remark: from the monotonicity property, the problem we are studying cannot present spurious eigenvalues

Uniform convergence

$$\|T - T_h\|_{\mathcal{L}(H,H)} \rightarrow 0$$

Convergence of eigenmodes (cont'ed)

Remark: from the monotonicity property, the problem we are studying cannot present spurious eigenvalues


Uniform convergence

$$\|T - T_h\|_{\mathcal{L}(H,H)} \rightarrow 0$$

Theorem

If T is selfadjoint and compact

Uniform convergence



Eigenmodes convergence

Convergence of eigenmodes (cont'ed)

Remark: from the monotonicity property, the problem we are studying cannot present spurious eigenvalues

Uniform convergence

$$\|T - T_h\|_{\mathcal{L}(H,H)} \rightarrow 0$$

Theorem

If T is selfadjoint and compact

Uniform convergence	\iff	Eigenmodes convergence
---------------------	--------	------------------------

Strategy

- 1) prove uniform convergence,
- 2) estimate the order of convergence

Galerkin approximation of compact operators

Bramble–Osborn '73

Osborn '75

Kolata '78

Galerkin approximation of compact operators

Bramble–Osborn '73

Osborn '75

Kolata '78

Céa's Lemma

$T_h = P_h T$, with P_h projection w.r.t. bilinear form a

Galerkin approximation of compact operators

Bramble–Osborn '73

Osborn '75

Kolata '78

Céa's Lemma

$T_h = P_h T$, with P_h projection w.r.t. bilinear form a

$$T - T_h = (I - P_h)T$$

If $I - P_h$ converges to zero **pointwise** and T is **compact**, then $T - T_h$ converges to zero **uniformly** (consequence of Banach–Steinhaus uniform boundedness theorem)

<Proof>

Estimating the order of convergence

Kolata '78

Babuška–Osborn '91

Estimating the order of convergence

If

Kolata '78

Babuška–Osborn '91

$$\begin{aligned}\|T - T_h\|_{\mathcal{L}(V,V)} &= \varepsilon(h) \\ \|T^* - T_h^*\|_{\mathcal{L}(V,V)} &= \varepsilon^*(h)\end{aligned}$$

then

$$\begin{aligned}|\lambda_i - \lambda_{i,h}| &\leq C(\varepsilon(h)\varepsilon^*(h))^{1/\alpha_i} \quad \alpha_i \text{ ascent of } \lambda_i I - T \\ \hat{\delta}(E_i, E_{i,h}) &\leq C\varepsilon(h)\end{aligned}$$

Estimating the order of convergence

If

Kolata '78

Babuška–Osborn '91

$$\begin{aligned}\|T - T_h\|_{\mathcal{L}(V,V)} &= \varepsilon(h) \\ \|T^* - T_h^*\|_{\mathcal{L}(V,V)} &= \varepsilon^*(h)\end{aligned}$$

then

$$\begin{aligned}|\lambda_i - \lambda_{i,h}| &\leq C(\varepsilon(h)\varepsilon^*(h))^{1/\alpha_i} \quad \alpha_i \text{ ascent of } \lambda_i I - T \\ \hat{\delta}(E_i, E_{i,h}) &\leq C\varepsilon(h)\end{aligned}$$

If T symmetric then

$$\begin{aligned}|\lambda_i - \lambda_{i,h}| &\leq C\varepsilon(h)^2 \\ \hat{\delta}(E_i, E_{i,h}) &\leq C\varepsilon(h)\end{aligned}$$

Piecewise linear FEs for Laplace eigenproblem

Simple proof that for $h < h_0$

$$\lambda_i \leq \lambda_{i,h} \leq \lambda_i + Ch^2$$

Piecewise linear FEs for Laplace eigenproblem

Simple proof that for $h < h_0$

$$\lambda_i \leq \lambda_{i,h} \leq \lambda_i + Ch^2$$

Take $E = \bigoplus_{k=1}^i E_k$, P_h elliptic projection, and $E_h = P_h E$.

Piecewise linear FEs for Laplace eigenproblem

Simple proof that for $h < h_0$

$$\lambda_i \leq \lambda_{i,h} \leq \lambda_i + Ch^2$$

Take $E = \bigoplus_{k=1}^i E_k$, P_h elliptic projection, and $E_h = P_h E$.
If h small enough, then $\dim E_h = i$

$$\lambda_{i,h} \leq \sup_{v \in E_h} \frac{\|\nabla v\|^2}{\|v\|^2}$$

Piecewise linear FEs for Laplace eigenproblem

Simple proof that for $h < h_0$

$$\lambda_i \leq \lambda_{i,h} \leq \lambda_i + Ch^2$$

Take $E = \bigoplus_{k=1}^i E_k$, P_h elliptic projection, and $E_h = P_h E$.
If h small enough, then $\dim E_h = i$

$$\lambda_{i,h} \leq \sup_{v \in E_h} \frac{\|\nabla v\|^2}{\|v\|^2} = \sup_{v \in E} \frac{\|\nabla P_h v\|^2}{\|P_h v\|^2}$$

Piecewise linear FEs for Laplace eigenproblem

Simple proof that for $h < h_0$

$$\lambda_i \leq \lambda_{i,h} \leq \lambda_i + Ch^2$$

Take $E = \bigoplus_{k=1}^i E_k$, P_h elliptic projection, and $E_h = P_h E$.
If h small enough, then $\dim E_h = i$

$$\lambda_{i,h} \leq \sup_{v \in E_h} \frac{\|\nabla v\|^2}{\|v\|^2} = \sup_{v \in E} \frac{\|\nabla P_h v\|^2}{\|P_h v\|^2} \leq \sup_{v \in E} \frac{\|\nabla v\|^2}{\|P_h v\|^2}$$

Piecewise linears for Laplace (cont'ed)

$$\lambda_{i,h} \leq \sup_{v \in E} \frac{\|\nabla v\|^2}{\|P_h v\|^2}$$

Piecewise linears for Laplace (cont'ed)

$$\lambda_{i,h} \leq \sup_{v \in E} \frac{\|\nabla v\|^2}{\|P_h v\|^2}$$

$$\|P_h v\| \geq \|v\| - \|P_h v - v\|$$

Piecewise linears for Laplace (cont'ed)

$$\lambda_{i,h} \leq \sup_{v \in E} \frac{\|\nabla v\|^2}{\|P_h v\|^2}$$

$$\begin{aligned}\|P_h v\| &\geq \|v\| - \|P_h v - v\| \\ \|P_h v - v\| &\leq Ch^2 \|\Delta v\|\end{aligned}$$

Piecewise linears for Laplace (cont'ed)

$$\lambda_{i,h} \leq \sup_{v \in E} \frac{\|\nabla v\|^2}{\|P_h v\|^2}$$

$$\begin{aligned}\|P_h v\| &\geq \|v\| - \|P_h v - v\| \\ \|P_h v - v\| &\leq Ch^2 \|\Delta v\| \leq Ch^2 \lambda_i \|v\|\end{aligned}$$

Piecewise linears for Laplace (cont'ed)

$$\lambda_{i,h} \leq \sup_{v \in E} \frac{\|\nabla v\|^2}{\|P_h v\|^2}$$

$$\begin{aligned}\|P_h v\| &\geq \|v\| - \|P_h v - v\| \\ \|P_h v - v\| &\leq Ch^2 \|\Delta v\| \leq Ch^2 \lambda_i \|v\| \\ \|P_h v\| &\geq \|v\| (1 - Ch^2)\end{aligned}$$

Piecewise linears for Laplace (cont'ed)

$$\lambda_{i,h} \leq \sup_{v \in E} \frac{\|\nabla v\|^2}{\|P_h v\|^2}$$

$$\begin{aligned}\|P_h v\| &\geq \|v\| - \|P_h v - v\| \\ \|P_h v - v\| &\leq Ch^2 \|\Delta v\| \leq Ch^2 \lambda_i \|v\| \\ \|P_h v\| &\geq \|v\| (1 - Ch^2)\end{aligned}$$

$$\lambda_{i,h} \leq \sup_{v \in E} \frac{\|\nabla v\|^2}{\|v\|^2} (1 + Ch^2) \leq \lambda_i + Ch^2$$

Computed eigenvalues: $[0, \pi]$ 1D interval, piecewise linear FEs

	$n = 8$	$n = 16$	$n = 32$
1	1.0129160450588	1.0032168743567	1.0008034482562
4	4.2095474481529	4.0516641802355	4.0128674974272
9	10.0802909335883	9.2631305555446	9.0652448637285
16	19.4536672593288	16.8381897926118	16.2066567209423
25	33.2628304890884	27.0649225609802	25.5059230069702

Computed eigenvalues: $[0, \pi]$ 1D interval, piecewise linear FEs

	$n = 8$	$n = 16$	$n = 32$
1	1.0129160450588	1.0032168743567	1.0008034482562
4	4.2095474481529	4.0516641802355	4.0128674974272
9	10.0802909335883	9.2631305555446	9.0652448637285
16	19.4536672593288	16.8381897926118	16.2066567209423
25	33.2628304890884	27.0649225609802	25.5059230069702

	$n = 64$	$n = 128$	$n = 256$
1	1.0002008137390	1.0000502004122	1.0000125499161
4	4.0032137930241	4.0008032549556	4.0002008016414
9	9.0162763381719	9.0040668861371	9.0010165838380
16	16.0514699897078	16.0128551720960	16.0032130198251
25	25.1257489536113	25.0313903532369	25.0078446408520

Conclusions on standard Galerkin approximation

Any finite element choice which provides a (pointwise) convergent scheme for the approximation of a problem with compact resolvent can be successfully applied to the approximation of the corresponding eigenvalue problem (uniform convergence).

No extra compatibility!

The curl curl operator

Let's consider the time harmonic model for Maxwell's system

$$\left(\varepsilon \frac{\partial}{\partial t} + \sigma \right) \mathcal{E} - \operatorname{curl} \mathcal{H} = \mathcal{J}, \quad \mu \frac{\partial}{\partial t} \mathcal{H} + \operatorname{curl} \mathcal{E} = 0.$$

The curl curl operator

Let's consider the time harmonic model for Maxwell's system

$$\left(\varepsilon \frac{\partial}{\partial t} + \sigma\right) \mathcal{E} - \operatorname{curl} \mathcal{H} = \mathcal{J}, \quad \mu \frac{\partial}{\partial t} \mathcal{H} + \operatorname{curl} \mathcal{E} = 0.$$

Time-harmonic model

$$\mathcal{E}(x, t) = e^{-i\omega t} E(x), \quad \mathcal{H}(x, t) = e^{-i\omega t} H(x)$$

$$-i\omega \left(\varepsilon + i\frac{\sigma}{\omega}\right) E - \operatorname{curl} H = J, \quad -i\omega \mu H + \operatorname{curl} E = 0.$$

Eliminating H

$$\operatorname{curl} (\mu^{-1} \operatorname{curl} E) - \omega^2 \left(\varepsilon + i\frac{\sigma}{\omega}\right) E = -i\omega J$$

The curl curl operator (cont'ed)

Change of notation $\mathbf{u} = E$, $\mathbf{f} = -i\omega J$

+ $\sigma = 0$

+ B.C.

$$\begin{cases} \operatorname{curl} (\mu^{-1} \operatorname{curl} \mathbf{u}) - \omega^2 \varepsilon \mathbf{u} = \mathbf{f} & \text{in } \Omega \\ \mathbf{u} \times \mathbf{n} = 0 & \text{on } \partial\Omega \end{cases}$$

The curl curl operator (cont'ed)

Change of notation $\mathbf{u} = E$, $\mathbf{f} = -i\omega J$

+ $\sigma = 0$

+ B.C.

$$\begin{cases} \operatorname{curl} (\mu^{-1} \operatorname{curl} \mathbf{u}) - \omega^2 \varepsilon \mathbf{u} = \mathbf{f} & \text{in } \Omega \\ \mathbf{u} \times \mathbf{n} = 0 & \text{on } \partial\Omega \end{cases}$$

The problem is well-posed if ω is not an interior Maxwell eigenvalue (otherwise $LHS = 0$ for \mathbf{u} eigenfunction)

The curl curl operator (cont'ed)

Variational formulation...

$V = H_0(\text{curl}; \Omega)$, find $\mathbf{u} \in V$ s.t.

$$(\mu^{-1} \text{curl } \mathbf{u}, \text{curl } \mathbf{v}) - \omega^2(\varepsilon \mathbf{u}, \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

The curl curl operator (cont'ed)

Variational formulation...

$V = H_0(\text{curl}; \Omega)$, find $\mathbf{u} \in V$ s.t.

$$(\mu^{-1} \text{curl } \mathbf{u}, \text{curl } \mathbf{v}) - \omega^2(\varepsilon \mathbf{u}, \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

Note: $\text{div } \mathbf{f} = 0$ implies $\text{div } \varepsilon \mathbf{u} = 0$

The curl curl operator (cont'ed)

Variational formulation...

$V = H_0(\text{curl}; \Omega)$, find $\mathbf{u} \in V$ s.t.

$$(\mu^{-1} \text{curl } \mathbf{u}, \text{curl } \mathbf{v}) - \omega^2(\varepsilon \mathbf{u}, \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

Note: $\text{div } \mathbf{f} = 0$ implies $\text{div } \varepsilon \mathbf{u} = 0$

...and its numerical approximation

$V_h \subset V$, find $\mathbf{u}_h \in V_h$ s.t.

$$(\mu^{-1} \text{curl } \mathbf{u}_h, \text{curl } \mathbf{v}) - \omega^2(\varepsilon \mathbf{u}_h, \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in V_h$$

The curl curl operator (cont'ed)

Time-harmonic Maxwell:

$$(\mu^{-1} \operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) - \omega^2 (\varepsilon \mathbf{u}, \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

The curl curl operator (cont'ed)

Time-harmonic Maxwell:

$$(\mu^{-1} \operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) - \omega^2(\varepsilon \mathbf{u}, \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

Maxwell interior eigenvalues:

$$(\mu^{-1} \operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) = \lambda^2(\varepsilon \mathbf{u}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

The curl curl operator (cont'ed)

Time-harmonic Maxwell:

$$(\mu^{-1} \operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) - \omega^2(\varepsilon \mathbf{u}, \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

Maxwell interior eigenvalues:

$$(\mu^{-1} \operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) = \lambda^2(\varepsilon \mathbf{u}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

The good approximation of the eigenvalues is clearly necessary for the well-posedness of the approximation to the time harmonic equations.

The curl curl operator (cont'ed)

Time-harmonic Maxwell:

$$(\mu^{-1} \operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) - \omega^2(\varepsilon \mathbf{u}, \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

Maxwell interior eigenvalues:

$$(\mu^{-1} \operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) = \lambda^2(\varepsilon \mathbf{u}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

The good approximation of the eigenvalues is clearly necessary for the well-posedness of the approximation to the time harmonic equations.

Indeed, it might be proved that this is also sufficient

The curl curl operator (cont'ed)

Time-harmonic Maxwell:

$$(\mu^{-1} \operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) - \omega^2(\varepsilon \mathbf{u}, \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

Maxwell interior eigenvalues:

$$(\mu^{-1} \operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) = \lambda^2(\varepsilon \mathbf{u}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$


The good approximation of the eigenvalues is clearly necessary for the well-posedness of the approximation to the time harmonic equations.

Indeed, it might be proved that this is also sufficient

N.B.: from now on we take $\mu = \varepsilon = 1$

2D numerical tests (curl curl operator)

Piecewise linear FEs on general meshes



The curl curl operator (cont'ed)

What is the difference with respect to previous abstract setting?

The curl curl operator (cont'ed)

What is the difference with respect to previous abstract setting?

$H = L^2$, $V = H_0(\text{curl}) \cap H(\text{div}^0)$, inclusion is compact

The curl curl operator (cont'ed)

What is the difference with respect to previous abstract setting?

$H = L^2$, $V = H_0(\text{curl}) \cap H(\text{div}^0)$, inclusion is compact

$a(\mathbf{u}, \mathbf{v}) = (\text{curl } \mathbf{u}, \text{curl } \mathbf{v})$ is coercive on V (Friedrichs inequality)

The curl curl operator (cont'ed)

What is the difference with respect to previous abstract setting?

$H = L^2$, $V = H_0(\text{curl}) \cap H(\text{div}^0)$, inclusion is compact

$a(\mathbf{u}, \mathbf{v}) = (\text{curl } \mathbf{u}, \text{curl } \mathbf{v})$ is coercive on V (Friedrichs inequality)

But. . . in this case V_h is not contained in V (divergence free constraint)

The curl curl operator (cont'ed)

What is the difference with respect to previous abstract setting?

$H = L^2$, $V = H_0(\text{curl}) \cap H(\text{div}^0)$, inclusion is compact

$a(\mathbf{u}, \mathbf{v}) = (\text{curl } \mathbf{u}, \text{curl } \mathbf{v})$ is coercive on V (Friedrichs inequality)

But. . . in this case V_h is not contained in V (divergence free constraint)

At least our theory was not wrong. . .

The curl curl operator (cont'ed)

Let's drop the divergence free constraint in the definition of our problem and see what happens

The curl curl operator (cont'ed)

Let's drop the divergence free constraint in the definition of our problem and see what happens

$$V = H_0(\text{curl})$$

Find $\lambda \in \mathbb{R}$ such that for some $\mathbf{u} \in V$ with $\mathbf{u} \neq 0$ it holds

$$(\text{curl } \mathbf{u}, \text{curl } \mathbf{v}) = \lambda(\mathbf{u}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

The curl curl operator (cont'ed)

Let's drop the divergence free constraint in the definition of our problem and see what happens

$$V = H_0(\text{curl})$$

Find $\lambda \in \mathbb{R}$ such that for some $\mathbf{u} \in V$ with $\mathbf{u} \neq 0$ it holds

$$(\text{curl } \mathbf{u}, \text{curl } \mathbf{v}) = \lambda(\mathbf{u}, \mathbf{v}) \quad \forall \mathbf{v} \in V$$

Remark: positive eigenvalues are the same as before. Eigenvalue $\lambda = 0$ is added to the spectrum with infinite dimensional eigenspace ∇H_0^1 . Consequence of Helmholtz decomposition which implies

$$V = [H_0(\text{curl}) \cap H(\text{div}^0)] \oplus \nabla H_0^1$$

The curl curl operator (cont'ed)

Let's try to use unconstrained formulation and consider finite element spaces which mimic at discrete level the Helmholtz decomposition.


The curl curl operator (cont'ed)

Let's try to use unconstrained formulation and consider finite element spaces which mimic at discrete level the Helmholtz decomposition. In particular, we want *gradients* to be well represented (so that zero frequency is well approximated).

The curl curl operator (cont'ed)

Let's try to use unconstrained formulation and consider finite element spaces which mimic at discrete level the Helmholtz decomposition. In particular, we want *gradients* to be well represented (so that zero frequency is well approximated).

Piecewise linear elements on criss-cross mesh have this property.




Results on criss-cross mesh


Mode		$n = 8$	$n = 16$	$n = 32$
(1,0)	1	1.00428	1.00107	1.00027
(0,1)	1	1.00428	1.00107	1.00027
(1,1)	2	2.01711	2.00428	2.00107
(2,0)	4	4.06804	4.01710	4.00428
(0,2)	4	4.06804	4.01710	4.00428
(2,1)	5	5.10634	5.02674	5.00669
(1,2)	5	5.10634	5.02674	5.00669
??	6	5.92293	5.98074	5.99518
(2,2)	8	8.27128	8.06845	8.01713
(3,0)	9	9.34085	9.08640	9.02166
(0,3)	9	9.34085	9.08640	9.02166
# zeros		63	255	1023

Spurious eigenmodes


1st spurious eigenfunction




2nd spurious eigenfunction




3rd spurious eigenfunction




4th spurious eigenfunction




Numerical results: edge elements




1st eigenfunction




2nd eigenfunction



3rd eigenfunction



4th eigenfunction



Numerical results: edge elements (cont'ed)

Mode		$n = 8$	$n = 16$	$n = 32$
(1,0)	1	0.99232	0.99806	0.99951
(0,1)	1	0.99914	0.99979	0.99994
(1,1)	2	2.00823	2.00212	2.00053
(2,0)	4	3.93162	3.98288	3.99572
(0,2)	4	3.93250	3.98294	3.99572
(2,1)	5	4.93116	4.98260	4.99564
(1,2)	5	5.05757	5.01511	5.00382
(2,2)	8	8.10159	8.03218	8.00844
(3,0)	9	8.62920	8.90607	8.97640
(0,3)	9	8.68245	8.92111	8.98027
# zeros		49	225	1023

Mixed formulations for curl curl eigenproblem

Kikuchi's formulation

Kikuchi '89

Find $\lambda \in \mathbb{R}$ s.t. for some $\mathbf{u} \in V$ with $\mathbf{u} \neq 0$ and $p \in H_0^1(\Omega) = Q$

$$(M_1) \quad \begin{aligned} (\operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) + (\mathbf{v}, \nabla p) &= \lambda(\mathbf{u}, \mathbf{v}) & \forall \mathbf{v} \in V \\ (\mathbf{u}, \nabla q) &= 0 & \forall q \in Q \end{aligned}$$

Mixed formulations for curl curl eigenproblem

Kikuchi's formulation

Kikuchi '89

Find $\lambda \in \mathbb{R}$ s.t. for some $\mathbf{u} \in V$ with $\mathbf{u} \neq 0$ and $p \in H_0^1(\Omega) = Q$

$$(M_1) \quad \begin{aligned} (\operatorname{curl} \mathbf{u}, \operatorname{curl} \mathbf{v}) + (\mathbf{v}, \nabla p) &= \lambda(\mathbf{u}, \mathbf{v}) & \forall \mathbf{v} \in V \\ (\mathbf{u}, \nabla q) &= 0 & \forall q \in Q \end{aligned}$$

Alternative mixed formulation B.–Fernandes–Gastaldi–Perugia '99

Find $\lambda \in \mathbb{R}$ s.t. for some $u \in \operatorname{curl} V = U$ with $u \neq 0$ and $\sigma \in V$

$$(M_2) \quad \begin{aligned} (\sigma, \tau) + (\operatorname{curl} \tau, u) &= 0 & \forall \tau \in V & \quad \sigma = -\operatorname{curl} u \\ (\operatorname{curl} \sigma, v) &= -\lambda(u, v) & \forall v \in U & \quad \operatorname{curl} \sigma = -\lambda u \end{aligned}$$

Discrete mixed formulations

Kikuchi's formulation

Find $\lambda_h \in \mathbb{R}$ s.t. for some $\mathbf{u}_h \in V_h$ with $\mathbf{u}_h \not\equiv 0$ and $p_h \in Q_h$

$$(M_{1,h}) \quad \begin{aligned} (\operatorname{curl} \mathbf{u}_h, \operatorname{curl} \mathbf{v}) + (\mathbf{v}, \nabla p_h) &= \lambda_h (\mathbf{u}_h, \mathbf{v}) & \forall \mathbf{v} \in V_h \\ (\mathbf{u}_h, \nabla q) &= 0 & \forall q \in Q_h \end{aligned}$$

Discrete mixed formulations

Kikuchi's formulation

Find $\lambda_h \in \mathbb{R}$ s.t. for some $\mathbf{u}_h \in V_h$ with $\mathbf{u}_h \not\equiv 0$ and $p_h \in Q_h$

$$(M_{1,h}) \quad \begin{aligned} (\operatorname{curl} \mathbf{u}_h, \operatorname{curl} \mathbf{v}) + (\mathbf{v}, \nabla p_h) &= \lambda_h (\mathbf{u}_h, \mathbf{v}) & \forall \mathbf{v} \in V_h \\ (\mathbf{u}_h, \nabla q) &= 0 & \forall q \in Q_h \end{aligned}$$

Alternative mixed formulation

Find $\lambda_h \in \mathbb{R}$ s.t. for some $u_h \in U_h$ with $u_h \not\equiv 0$ and $\sigma_h \in \Sigma_h$

$$(M_{2,h}) \quad \begin{aligned} (\sigma_h, \tau) + (\operatorname{curl} \tau, u_h) &= 0 & \forall \tau \in \Sigma_h \\ (\operatorname{curl} \sigma_h, v) &= -\lambda_h (u_h, v) & \forall v \in U_h \end{aligned}$$

Discrete mixed formulations

Kikuchi's formulation

Find $\lambda_h \in \mathbb{R}$ s.t. for some $\mathbf{u}_h \in V_h$ with $\mathbf{u}_h \not\equiv 0$ and $p_h \in Q_h$

$$(M_{1,h}) \quad \begin{aligned} (\operatorname{curl} \mathbf{u}_h, \operatorname{curl} \mathbf{v}) + (\mathbf{v}, \nabla p_h) &= \lambda_h (\mathbf{u}_h, \mathbf{v}) & \forall \mathbf{v} \in V_h \\ (\mathbf{u}_h, \nabla q) &= 0 & \forall q \in Q_h \end{aligned}$$

Alternative mixed formulation

Find $\lambda_h \in \mathbb{R}$ s.t. for some $u_h \in U_h$ with $u_h \not\equiv 0$ and $\sigma_h \in \Sigma_h$

$$(M_{2,h}) \quad \begin{aligned} (\sigma_h, \tau) + (\operatorname{curl} \tau, u_h) &= 0 & \forall \tau \in \Sigma_h \\ (\operatorname{curl} \sigma_h, v) &= -\lambda_h (u_h, v) & \forall v \in U_h \end{aligned}$$

Theorem

$M_{1,h} \iff M_{2,h}$ and equiv. to original problem ($\lambda_h \neq 0$) provided

$$\nabla Q_h \subset V_h \text{ and } \operatorname{curl} V_h \subset U_h$$

In 2D this is simply *rotated* mixed Laplace

Find $\lambda_h \in \mathbb{R}$ s.t. for some $u_h \in U_h$ with $u_h \neq 0$ and $\sigma_h \in \Sigma_h$

$$(M_{2,h}) \quad \begin{aligned} (\sigma_h, \tau) + (\operatorname{curl} \tau, u_h) &= 0 \quad \forall \tau \in \Sigma_h \\ (\operatorname{curl} \sigma_h, v) &= -\lambda_h (u_h, v) \quad \forall v \in U_h \end{aligned}$$

In 2D this is simply *rotated* mixed Laplace

Find $\lambda_h \in \mathbb{R}$ s.t. for some $u_h \in U_h$ with $u_h \neq 0$ and $\sigma_h \in \Sigma_h$

$$(M_{2,h}) \quad \begin{aligned} (\sigma_h, \tau) + (\operatorname{curl} \tau, u_h) &= 0 \quad \forall \tau \in \Sigma_h \\ (\operatorname{curl} \sigma_h, v) &= -\lambda_h(u_h, v) \quad \forall v \in U_h \end{aligned}$$

$$\begin{aligned} (\sigma_h, \tau) + (\operatorname{div} \tau, u_h) &= 0 \quad \forall \tau \in \Sigma_h \\ (\operatorname{div} \sigma_h, v) &= -\lambda_h(u_h, v) \quad \forall v \in U_h \end{aligned}$$

Eigenproblems in mixed form

Mercier–Osborn–Rappaz–Raviart '81
B.–Brezzi–Gastaldi '97-'00

Eigenproblems in mixed form

Mercier–Osborn–Rappaz–Raviart '81
B.–Brezzi–Gastaldi '97-'00

Let's start with Laplace problem $-\Delta u = g$

Source problem

$$\sigma \in H(\operatorname{div}; \Omega) = \Sigma, \quad u \in L^2(\Omega) = U$$

$$\begin{cases} (\sigma, \tau) + (\operatorname{div} \tau, u) = 0 & \forall \tau \in \Sigma & (\sigma = \nabla u) \\ (\operatorname{div} \sigma, v) = -(g, v) & \forall v \in U & (-\operatorname{div} \sigma = g) \end{cases}$$

Eigenproblems in mixed form

Mercier–Osborn–Rappaz–Raviart '81
B.–Brezzi–Gastaldi '97-'00

Let's start with Laplace problem $-\Delta u = g$

Source problem

$$\sigma \in H(\operatorname{div}; \Omega) = \Sigma, \quad u \in L^2(\Omega) = U$$

$$\begin{cases} (\sigma, \tau) + (\operatorname{div} \tau, u) = 0 & \forall \tau \in \Sigma & (\sigma = \nabla u) \\ (\operatorname{div} \sigma, v) = -(g, v) & \forall v \in U & (-\operatorname{div} \sigma = g) \end{cases}$$

Matrix form ($\Sigma_h \subset \Sigma, U_h \subset U$)

$$\begin{pmatrix} A & B^t \\ B & 0 \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix} = \begin{pmatrix} 0 \\ -\tilde{g} \end{pmatrix}$$

Laplace eigenproblem in mixed form

Find $\lambda \in \mathbb{R}$ such that for some $(\sigma, u) \in \Sigma \times U$ with $u \neq 0$

$$\begin{cases} (\sigma, \tau) + (\operatorname{div} \tau, u) = 0 & \forall \tau \in \Sigma \\ (\operatorname{div} \sigma, v) = -\lambda(u, v) & \forall v \in U \end{cases} \quad (-\Delta u = \lambda u)$$

Laplace eigenproblem in mixed form

Find $\lambda \in \mathbb{R}$ such that for some $(\sigma, u) \in \Sigma \times U$ with $u \neq 0$

$$\begin{cases} (\sigma, \tau) + (\operatorname{div} \tau, u) = 0 & \forall \tau \in \Sigma \\ (\operatorname{div} \sigma, v) = -\lambda(u, v) & \forall v \in U \end{cases} \quad (-\Delta u = \lambda u)$$

Matrix form ($\Sigma_h \subset \Sigma$, $U_h \subset U$)

$$\begin{pmatrix} A & B^t \\ B & 0 \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix} = -\tilde{\lambda} \begin{pmatrix} 0 & 0 \\ 0 & M_U \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix}$$

Laplace eigenproblem in mixed form

Find $\lambda \in \mathbb{R}$ such that for some $(\sigma, u) \in \Sigma \times U$ with $u \neq 0$

$$\begin{cases} (\sigma, \tau) + (\operatorname{div} \tau, u) = 0 & \forall \tau \in \Sigma \\ (\operatorname{div} \sigma, v) = -\lambda(u, v) & \forall v \in U \end{cases} \quad (-\Delta u = \lambda u)$$

Matrix form ($\Sigma_h \subset \Sigma$, $U_h \subset U$)

$$\begin{pmatrix} A & B^t \\ B & 0 \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix} = -\tilde{\lambda} \begin{pmatrix} 0 & 0 \\ 0 & M_U \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix}$$

Remark. Similarly, one can deal with problems of the type

$$\begin{pmatrix} A & B^t \\ B & 0 \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix} = \lambda \begin{pmatrix} M_\Sigma & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix}$$

Definition of the resolvent operator

A first natural (but wrong) definition

$$T_1 : U \rightarrow \Sigma \times U$$

$$T_1(g) = (\sigma, u)$$

One would like to compute eigenvalues. . .

Definition of the resolvent operator

A first natural (but wrong) definition

$$T_1 : U \rightarrow \Sigma \times U$$

$$T_1(g) = (\sigma, u)$$

One would like to compute eigenvalues. . .

$$T_2 : (\Sigma \times U)' \rightarrow \Sigma \times U$$

$$T_2(f, g) = (\sigma, u) \text{ with}$$

$$\begin{cases} (\sigma, \tau) + (\operatorname{div} \tau, u) = \langle f, v \rangle & \forall \tau \in \Sigma \\ (\operatorname{div} \sigma, v) = -(g, v) & \forall v \in U \end{cases}$$

Definition of the resolvent operator

A first natural (but wrong) definition

$$T_1 : U \rightarrow \Sigma \times U$$

$$T_1(g) = (\sigma, u)$$

One would like to compute eigenvalues. . .

$$T_2 : (\Sigma \times U)' \rightarrow \Sigma \times U$$

$$T_2(f, g) = (\sigma, u) \text{ with}$$

$$\begin{cases} (\sigma, \tau) + (\operatorname{div} \tau, u) = \langle f, v \rangle & \forall \tau \in \Sigma \\ (\operatorname{div} \sigma, v) = -(g, v) & \forall v \in U \end{cases}$$

$$T_{\Sigma U} \boxed{\begin{array}{ccc} (f, g) & \xrightarrow{\text{cutoff}} & (0, g) \xrightarrow{T_2} (\sigma, u) \\ L^2 \times L^2 & \longrightarrow & L^2 \times L^2 \end{array}} \text{ is compact}$$

Uniform convergence?

Let's try to follow Kolata's argument

$$T_{\Sigma U} - T_{\Sigma U, h} = (I - Q_h)T_{\Sigma U}$$

Uniform convergence?

Let's try to follow Kolata's argument

$$T_{\Sigma U} - T_{\Sigma U, h} = (I - Q_h)T_{\Sigma U}$$

✓ $\|(I - Q_h)(\sigma, u)\|_{\Sigma \times U} \rightarrow 0$ for all $(\sigma, u) \in \Sigma \times U$

Uniform convergence?

Let's try to follow Kolata's argument

$$T_{\Sigma U} - T_{\Sigma U, h} = (I - Q_h)T_{\Sigma U}$$

✓ $\|(I - Q_h)(\sigma, u)\|_{\Sigma \times U} \rightarrow 0$ for all $(\sigma, u) \in \Sigma \times U$

✗ $T_{\Sigma U} : L^2 \times L^2 \rightarrow \Sigma \times U$ is not compact

Uniform convergence?

Let's try to follow Kolata's argument

$$T_{\Sigma U} - T_{\Sigma U, h} = (I - Q_h)T_{\Sigma U}$$

✓ $\|(I - Q_h)(\sigma, u)\|_{\Sigma \times U} \rightarrow 0$ for all $(\sigma, u) \in \Sigma \times U$

✗ $T_{\Sigma U} : L^2 \times L^2 \rightarrow \Sigma \times U$ is not compact

✗ $T_{\Sigma U} : \Sigma \times U \rightarrow \Sigma \times U$ is not compact either

Uniform convergence?

Let's try to follow Kolata's argument

$$T_{\Sigma U} - T_{\Sigma U, h} = (I - Q_h)T_{\Sigma U}$$

✓ $\|(I - Q_h)(\sigma, u)\|_{\Sigma \times U} \rightarrow 0$ for all $(\sigma, u) \in \Sigma \times U$

✗ $T_{\Sigma U} : L^2 \times L^2 \rightarrow \Sigma \times U$ is not compact

✗ $T_{\Sigma U} : \Sigma \times U \rightarrow \Sigma \times U$ is not compact either

Remark: standard mixed estimates don't help

$$\|\sigma - \sigma_h\|_{\Sigma} + \|u - u_h\|_U \leq C \inf_{\tau_h, v_h} \left(\underbrace{\|\sigma - \tau_h\|_{\Sigma}}_{O(1)} + \underbrace{\|u - v_h\|_U}_{O(h)} \right)$$

Definition of the resolvent operator (cont'ed)

A better definition

$$T_U : U \rightarrow U$$

$\sigma \in \Sigma$, $T_U g \in U$ such that

$$\begin{cases} (\sigma, \tau) + (\operatorname{div} \tau, T_U g) = 0 & \forall \tau \in \Sigma \\ (\operatorname{div} \sigma, v) = -(g, v) & \forall v \in U \end{cases}$$

Definition of the resolvent operator (cont'ed)

A better definition

$$T_U : U \rightarrow U$$

$\sigma \in \Sigma$, $T_U g \in U$ such that

$$\begin{cases} (\sigma, \tau) + (\operatorname{div} \tau, T_U g) = 0 & \forall \tau \in \Sigma \\ (\operatorname{div} \sigma, v) = -(g, v) & \forall v \in U \end{cases}$$

Remark: operator is now compact, but standard mixed estimates don't help again

$$\|\sigma - \sigma_h\|_{\Sigma} + \|u - u_h\|_U \leq C \inf_{\tau_h, v_h} \left(\underbrace{\|\sigma - \tau_h\|_{\Sigma}}_{O(1)} + \underbrace{\|u - v_h\|_U}_{O(h)} \right)$$

Definition of the resolvent operator (cont'ed)

A better definition

$$T_U : U \rightarrow U$$

$\sigma \in \Sigma$, $T_U g \in U$ such that

$$\begin{cases} (\sigma, \tau) + (\operatorname{div} \tau, T_U g) = 0 & \forall \tau \in \Sigma \\ (\operatorname{div} \sigma, v) = -(g, v) & \forall v \in U \end{cases}$$

Remark: operator is now compact, but standard mixed estimates don't help again

$$||\sigma - \sigma_h||_{\Sigma} + ||u - u_h||_U \leq C \inf_{\tau_h, v_h} \left(\underbrace{||\sigma - \tau_h||_{\Sigma}}_{O(1)} + \underbrace{||u - v_h||_U}_{O(h)} \right)$$

Remark: we need an estimate for u_h which does not involve $\operatorname{div} \sigma$

Uniform convergence $\|T_U - T_{U,h}\| \rightarrow 0$

► Ellipticity in the kernel

$$\|\tau_h\|_{L^2}^2 \geq \alpha \|\tau_h\|_{\Sigma}^2 \text{ for all } \tau_h \in \Sigma_h \text{ s.t. } \{(\operatorname{div} \tau_h, v) = 0, \forall v \in U_h\}$$

Uniform convergence $\|T_U - T_{U,h}\| \rightarrow 0$

- Ellipticity in the kernel

$$\|\tau_h\|_{L^2}^2 \geq \alpha \|\tau_h\|_{\Sigma}^2 \text{ for all } \tau_h \in \Sigma_h \text{ s.t. } \{(\operatorname{div} \tau_h, v) = 0, \forall v \in U_h\}$$

- Fortin operator $\Pi_h : \Sigma^+ \rightarrow \Sigma_h$ s.t.

$$(\operatorname{div}(\sigma - \Pi_h \sigma), v) = 0 \quad \forall v \in U_h$$

$$\|\Pi_h \sigma\|_{\Sigma} \leq C \|\sigma\|_{\Sigma^+}$$

Uniform convergence $\|T_U - T_{U,h}\| \rightarrow 0$

- Ellipticity in the kernel

$$\|\tau_h\|_{L^2}^2 \geq \alpha \|\tau_h\|_{\Sigma}^2 \text{ for all } \tau_h \in \Sigma_h \text{ s.t. } \{(\operatorname{div} \tau_h, v) = 0, \forall v \in U_h\}$$

- Fortin operator $\Pi_h : \Sigma^+ \rightarrow \Sigma_h$ s.t.

$$(\operatorname{div}(\sigma - \Pi_h \sigma), v) = 0 \quad \forall v \in U_h$$

$$\|\Pi_h \sigma\|_{\Sigma} \leq C \|\sigma\|_{\Sigma^+}$$

$$\|\sigma - \sigma_h\|_{L^2} \leq C \left(\|\sigma - \Pi_h \sigma\|_{L^2} + (1/\sqrt{\alpha}) \inf_{v_h \in U_h} \|u - v_h\|_U \right)$$

$$\|u - u_h\|_U \leq C \left(\inf_{v_h \in U_h} \|u - v_h\|_U + \|\sigma - \sigma_h\|_{L^2} \right)$$

<Proof>

Fortid condition

Definition

The spaces Σ_h , U_h satisfy the Fortid condition if there exists a Fortin operator which converges strongly to the identity operator, namely

$\Pi_h : \Sigma^+ \rightarrow \Sigma_h$ s.t.

$$(\operatorname{div}(\sigma - \Pi_h \sigma), v) = 0 \quad \forall v \in U_h$$

$$\|\Pi_h \sigma\|_{\Sigma} \leq C \|\sigma\|_{\Sigma^+}$$

$$\|I - \Pi_h\|_{\mathcal{L}(\Sigma^+, L^2)} \rightarrow 0$$

Final convergence result

Theorem

Assume ellipticity in the kernel and Fortin condition

For any $N \in \mathbb{N}$ define $\rho_N(h) :]0, 1] \rightarrow \mathbb{R}$ as

$$\rho_N(h) = \sup_{\substack{u \in \bigoplus_{i=1}^{m(N)} E_i}} \left(\inf_{v_h} \|u - v_h\|_U + \|\nabla u - \Pi_h \nabla u\|_{L^2} \right)$$

Then $\|T_U - T_{U,h}\|_{\mathcal{L}(U,U)} \rightarrow 0$ and the following estimates hold true

$$\sum_{i=1}^{m(N)} |\lambda_i - \lambda_{i,h}| \leq C(\rho_N(h))^2$$

$$\hat{\delta} \left(\bigoplus_{i=1}^{m(N)} E_i, \bigoplus_{i=1}^{m(N)} E_{i,h} \right) \leq C \rho_N(h)$$

Back to the criss-cross example

Crisscross mesh

$$\Sigma_h = \{\text{continuous p.w. linears (componentwise)}\}$$

$$U_h = \operatorname{div} \Sigma_h \subset \{\text{p.w. constants}\}$$

Back to the criss-cross example

Crisscross mesh

$\Sigma_h = \{\text{continuous p.w. linears (componentwise)}\}$

$U_h = \text{div } \Sigma_h \subset \{\text{p.w. constants}\}$

Theorem

With the above choice of spaces, there exist a sequence $\{g_h\} \subset U$ with $\|g_h\|_0 = 1$ s.t.

$$\|u - u_h\|_U \not\rightarrow 0$$

that is $\|T_U - T_{U,h}\|_{\mathcal{L}(U,U)} \not\rightarrow 0$

Back to the criss-cross example

Crisscross mesh

$\Sigma_h = \{\text{continuous p.w. linears (componentwise)}\}$

$U_h = \text{div } \Sigma_h \subset \{\text{p.w. constants}\}$

Theorem

With the above choice of spaces, there exist a sequence $\{g_h\} \subset U$ with $\|g_h\|_0 = 1$ s.t.

$$\|u - u_h\|_U \not\rightarrow 0$$

that is $\|T_U - T_{U,h}\|_{\mathcal{L}(U,U)} \not\rightarrow 0$

Proof

Use inequality by Qin '94 based on idea of Boland–Nicolaides '85
<More detail>

A positive example

General mesh (triangles, parallelograms, tetrahedrons, parallelepipeds)

Σ_h : Raviart–Thomas space of order k

U_h : \mathcal{P}_{k-1} or tensor product polynomials \mathcal{Q}_{k-1}

A positive example

General mesh (triangles, parallelograms, tetrahedrons, parallelepipeds)

Σ_h : Raviart–Thomas space of order k

U_h : \mathcal{P}_{k-1} or tensor product polynomials \mathcal{Q}_{k-1}

Fortid: the interpolant is a Fortin operator.

See also Falk–Osborn '80

A positive example

General mesh (triangles, parallelograms, tetrahedrons, parallelepipeds)

Σ_h : Raviart–Thomas space of order k

U_h : \mathcal{P}_{k-1} or tensor product polynomials \mathcal{Q}_{k-1}

Fortid: the interpolant is a Fortin operator.

See also **Falk–Osborn '80**

Convergence: $O(h^{2k})$ eigenvalues, $O(h^k)$ eigenfunctions.

A positive example

General mesh (triangles, parallelograms, tetrahedrons, parallelepipeds)

Σ_h : Raviart–Thomas space of order k

U_h : \mathcal{P}_{k-1} or tensor product polynomials \mathcal{Q}_{k-1}

Fortid: the interpolant is a Fortin operator.

See also Falk–Osborn '80

Convergence: $O(h^{2k})$ eigenvalues, $O(h^k)$ eigenfunctions.

Remark: in 2D edge elements can be chosen as *rotated* Raviart–Thomas spaces

General theory

$\begin{pmatrix} 0 \\ g \end{pmatrix}$ -type problems

$$\begin{pmatrix} A & B^t \\ B & 0 \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix} = -\lambda \begin{pmatrix} 0 & 0 \\ 0 & M_U \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix}$$

Laplace, biharmonic, Maxwell (1), . . .

General theory

$\begin{pmatrix} 0 \\ g \end{pmatrix}$ -type problems

$$\begin{pmatrix} A & B^t \\ B & 0 \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix} = -\lambda \begin{pmatrix} 0 & 0 \\ 0 & M_U \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix}$$

Laplace, biharmonic, Maxwell (1), . . .

$\begin{pmatrix} f \\ 0 \end{pmatrix}$ -type problems

$$\begin{pmatrix} A & B^t \\ B & 0 \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix} = \lambda \begin{pmatrix} M_\Sigma & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \tilde{\sigma} \\ \tilde{u} \end{pmatrix}$$

Stokes, Laplace with Lagrange multipliers, Maxwell (2), . . .

<More detail>

Some References

- I. Babuška and J. Osborn, Eigenvalue problems, in Handbook of Numerical Analysis, P. Ciarlet and J. Lions, eds., vol. II, Elsevier Science Publishers B.V., North Holland, 1991, pp. 641-788.
- D. Boffi, Fortin operator and discrete compactness for edge elements, Numer. Math., 87 (2000), pp. 229-246.
- D. Boffi, A note on the de Rham complex and a discrete compactness property, Appl. Math. Lett., 14 (2001), pp. 33-38.
- D. Boffi, Compatible discretizations for eigenvalue problems.

Compatible spatial discretizations, 121-142, IMA Vol. Math. Appl., 142, Springer, New York, 2006.

- D. Boffi. Approximation of eigenvalues in mixed form, discrete compactness property, and application to hp mixed finite elements. Computer Methods in Applied Mechanics and Engineering, to appear.
- D. Boffi, F. Brezzi, and L. Gastaldi, On the convergence of eigenvalues for mixed formulations, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 25 (1997), pp. 131-154 (1998). Dedicated to Ennio De Giorgi.
- D. Boffi, F. Brezzi, and L. Gastaldi, On the problem of spurious eigenvalues in the approximation of linear elliptic problems in mixed form, Math. Comp., 69 (2000), pp. 121-140.

- D. Boffi, P. Fernandes, L. Gastaldi, and I. Perugia, Computational models of electromagnetic resonators: analysis of edge element approximation, SIAM J. Numer. Anal., 36 (1999), pp. 1264-1290.
- D. Boffi, F. Kikuchi, J. Schöberl, Edge element computation of Maxwell's eigenvalues on general quadrilateral meshes. Math. Models Methods Appl. Sci. 16 (2006), no. 2, 265-273.
- L. Demkowicz, P. Monk, L. Vardapetyan, W. Rachowicz, de Rham diagram for hp finite element spaces. Comput. Math. Appl. 39 (2000), no. 7-8, 29-38
- T. Kato, Perturbation theory for linear operators, Classics in Mathematics, Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition.

- W.G. Kolata, Approximation in variationally posed eigenvalue problems. Numer. Math. 29 (1977/78), no. 2, 159-171.
- B. Mercier, J. Osborn, J. Rappaz, P.-A. Raviart, Eigenvalue approximation by mixed and hybrid methods. Math. Comp. 36 (1981), no. 154, 427-453.
- P.-A. Raviart, J.-M. Thomas, Introduction à l'analyse numérique des équations aux dérivées partielles. Masson, Paris, 1983

Conclusions

- ➡ Standard Galerkin approximation of eigenvalue problems is straightforward extension of usual theory

Conclusions

- ➡ Standard Galerkin approximation of eigenvalue problems is straightforward extension of usual theory
- ➡ Curl curl operator has driven us into more complicated problems

Conclusions

- ➡ Standard Galerkin approximation of eigenvalue problems is straightforward extension of usual theory
- ➡ Curl curl operator has driven us into more complicated problems
- ➡ Mixed problems require special theory (two different theories for $\begin{pmatrix} f \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ g \end{pmatrix}$ problems)

Conclusions

- ➡ Standard Galerkin approximation of eigenvalue problems is straightforward extension of usual theory
- ➡ Curl curl operator has driven us into more complicated problems
- ➡ Mixed problems require special theory (two different theories for $\begin{pmatrix} f \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ g \end{pmatrix}$ problems)

Conclusions

- ➡ Standard Galerkin approximation of eigenvalue problems is straightforward extension of usual theory
- ➡ Curl curl operator has driven us into more complicated problems
- ➡ Mixed problems require special theory (two different theories for $\begin{pmatrix} f \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ g \end{pmatrix}$ problems)

Conclusions

- ➡ Standard Galerkin approximation of eigenvalue problems is straightforward extension of usual theory
- ➡ Curl curl operator has driven us into more complicated problems
- ➡ Mixed problems require special theory (two different theories for $\begin{pmatrix} f \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ g \end{pmatrix}$ problems)

Conclusions


- ➡ Standard Galerkin approximation of eigenvalue problems is straightforward extension of usual theory
- ➡ Curl curl operator has driven us into more complicated problems
- ➡ Mixed problems require special theory (two different theories for $\begin{pmatrix} f \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ g \end{pmatrix}$ problems)

Conclusions


- ➡ Standard Galerkin approximation of eigenvalue problems is straightforward extension of usual theory
- ➡ Curl curl operator has driven us into more complicated problems
- ➡ Mixed problems require special theory (two different theories for $\begin{pmatrix} f \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ g \end{pmatrix}$ problems)

The end


Examples of good or bad convergence



Good



Bad (missing)



Bad (spurious)

<Back>

$$\|T - T_h\|_{\mathcal{L}(H,H)} \rightarrow 0$$

$$\|(I - P_h)(u)\|_H \rightarrow 0 \text{ for any } u \in V$$

Banach-Steinhaus thm. implies $\|I - P_h\|_{\mathcal{L}(V,H)} \leq C$

$$\|T - T_h\|_{\mathcal{L}(H,H)} \rightarrow 0$$

$\|(I - P_h)(u)\|_H \rightarrow 0$ for any $u \in V$

Banach-Steinhaus thm. implies $\|I - P_h\|_{\mathcal{L}(V,H)} \leq C$

By contradiction $\|(I - P_h)T\|_{\mathcal{L}(H,H)} \not\rightarrow 0$

Then (up to subsequences) $\|(I - P_h)Tf_h\|_H \not\rightarrow 0$ with $\|f_h\|_H \leq 1$

$$\|T - T_h\|_{\mathcal{L}(H,H)} \rightarrow 0$$

$\|(I - P_h)(u)\|_H \rightarrow 0$ for any $u \in V$

Banach-Steinhaus thm. implies $\|I - P_h\|_{\mathcal{L}(V,H)} \leq C$

By contradiction $\|(I - P_h)T\|_{\mathcal{L}(H,H)} \not\rightarrow 0$

Then (up to subsequences) $\|(I - P_h)Tf_h\|_H \not\rightarrow 0$ with $\|f_h\|_H \leq 1$

If $T : H \rightarrow V$ is compact then $Tf_h \rightarrow w$ in V

$$\|T - T_h\|_{\mathcal{L}(H,H)} \rightarrow 0$$

$\|(I - P_h)(u)\|_H \rightarrow 0$ for any $u \in V$

Banach-Steinhaus thm. implies $\|I - P_h\|_{\mathcal{L}(V,H)} \leq C$

By contradiction $\|(I - P_h)T\|_{\mathcal{L}(H,H)} \not\rightarrow 0$

Then (up to subsequences) $\|(I - P_h)Tf_h\|_H \not\rightarrow 0$ with $\|f_h\|_H \leq 1$

If $T : H \rightarrow V$ is compact then $Tf_h \rightarrow w$ in V

Take h small enough s.t. $\|Tf_h - w\|_V + \|(I - P_h)w\|_H \leq \varepsilon$

$$\|T - T_h\|_{\mathcal{L}(H,H)} \rightarrow 0$$

$$\|(I - P_h)(u)\|_H \rightarrow 0 \text{ for any } u \in V$$

Banach-Steinhaus thm. implies $\|I - P_h\|_{\mathcal{L}(V,H)} \leq C$

By contradiction $\|(I - P_h)T\|_{\mathcal{L}(H,H)} \not\rightarrow 0$

Then (up to subsequences) $\|(I - P_h)Tf_h\|_H \not\rightarrow 0$ with $\|f_h\|_H \leq 1$

If $T : H \rightarrow V$ is compact then $Tf_h \rightarrow w$ in V

Take h small enough s.t. $\|Tf_h - w\|_V + \|(I - P_h)w\|_H \leq \varepsilon$, then
 $\|(I - P_h)Tf_h\|_H \leq \|(I - P_h)(Tf_h - w)\|_H + \|(I - P_h)w\|_H \leq (C + 1)\varepsilon$

$$\|T - T_h\|_{\mathcal{L}(H,H)} \rightarrow 0$$

$\|(I - P_h)(u)\|_H \rightarrow 0$ for any $u \in V$

Banach-Steinhaus thm. implies $\|I - P_h\|_{\mathcal{L}(V,H)} \leq C$

By contradiction $\|(I - P_h)T\|_{\mathcal{L}(H,H)} \not\rightarrow 0$

Then (up to subsequences) $\|(I - P_h)Tf_h\|_H \not\rightarrow 0$ with $\|f_h\|_H \leq 1$

If $T : H \rightarrow V$ is compact then $Tf_h \rightarrow w$ in V

Take h small enough s.t. $\|Tf_h - w\|_V + \|(I - P_h)w\|_H \leq \varepsilon$, then
 $\|(I - P_h)Tf_h\|_H \leq \|(I - P_h)(Tf_h - w)\|_H + \|(I - P_h)w\|_H \leq (C + 1)\varepsilon$

Remark. Same argument shows $\|T - T_h\|_{\mathcal{L}(V,V)} \rightarrow 0$ under the hypotheses $T : V \rightarrow V$ compact and $\|I - P_h\|_{\mathcal{L}(V,V)} \leq C$

$$||T - T_h||_{\mathcal{L}(H,H)} \rightarrow 0 \text{ (cont'ed)}$$

A direct proof using a priori estimates

$$||(T - T_h)f||_H = ||u - u_h||_H \leq Ch^\alpha ||f||_H$$

<Back>

$P = L^2$ -projection onto U_h

$$\begin{aligned} ||\Pi_h \sigma - \sigma_h||_{L^2}^2 &= (\Pi_h \sigma - \sigma, \Pi_h \sigma - \sigma_h) + (\sigma - \sigma_h, \Pi_h \sigma - \sigma_h) \\ &= (\Pi_h \sigma - \sigma, \Pi_h \sigma - \sigma_h) - (\operatorname{div}(\Pi_h \sigma - \sigma_h), u - Pu) \\ &\leq ||\Pi_h \sigma - \sigma||_{L^2} ||\Pi_h \sigma - \sigma_h||_{L^2} + ||\operatorname{div}(\Pi_h \sigma - \sigma_h)||_{L^2} ||u - Pu||_U \\ &\leq ||\Pi_h \sigma - \sigma_h||_{L^2} (||\Pi_h \sigma - \sigma||_{L^2} + (1/\sqrt{\alpha}) ||u - Pu||_U) \end{aligned}$$

$P = L^2$ -projection onto U_h

$$\begin{aligned}
 ||\Pi_h \sigma - \sigma_h||_{L^2}^2 &= (\Pi_h \sigma - \sigma, \Pi_h \sigma - \sigma_h) + (\sigma - \sigma_h, \Pi_h \sigma - \sigma_h) \\
 &= (\Pi_h \sigma - \sigma, \Pi_h \sigma - \sigma_h) - (\operatorname{div}(\Pi_h \sigma - \sigma_h), u - Pu) \\
 &\leq ||\Pi_h \sigma - \sigma||_{L^2} ||\Pi_h \sigma - \sigma_h||_{L^2} + ||\operatorname{div}(\Pi_h \sigma - \sigma_h)||_{L^2} ||u - Pu||_U \\
 &\leq ||\Pi_h \sigma - \sigma_h||_{L^2} (||\Pi_h \sigma - \sigma||_{L^2} + (1/\sqrt{\alpha}) ||u - Pu||_U)
 \end{aligned}$$

$$\begin{aligned}
 ||Pu - u_h||_U &\leq C \sup_{\tau_h} \frac{(Pu - u_h, \operatorname{div} \tau_h)}{||\tau_h||_\Sigma} \\
 &\leq C \sup_{\tau_h} \frac{(Pu - u, \operatorname{div} \tau_h) + (u - u_h, \operatorname{div} \tau_h)}{||\tau_h||_\Sigma} \\
 &\leq C \left(||Pu - u||_U + \sup_{\tau_h} \frac{-(\sigma - \sigma_h, \tau_h)}{||\tau_h||_\Sigma} \right) \\
 &\leq C (||Pu - u||_U + ||\sigma - \sigma_h||_{L^2})
 \end{aligned}$$

<Back>

Inf-sup condition for our problem

$$\inf_{v \in U_h} \sup_{\tau \in \Sigma_h} \frac{(\operatorname{div} \tau, v)}{\|\tau\|_{H(\operatorname{div})} \|v\|_{L^2}} \geq \beta > 0$$

Inf-sup condition for our problem

$$\inf_{v \in U_h} \sup_{\tau \in \Sigma_h} \frac{(\operatorname{div} \tau, v)}{\|\tau\|_{H(\operatorname{div})} \|v\|_{L^2}} \geq \beta > 0$$

cannot be improved: there exists a sequence (properly chosen linear combination of checkerboards on the macroelements) $\{\tilde{v}_h\}$ such that

$$|(\operatorname{div} \tau, \tilde{v}_h)| \leq C \|\tau\|_{L^2} \|\tilde{v}_h\|_{L^2}$$

Inf-sup condition for our problem

$$\inf_{v \in U_h} \sup_{\tau \in \Sigma_h} \frac{(\operatorname{div} \tau, v)}{\|\tau\|_{H(\operatorname{div})} \|v\|_{L^2}} \geq \beta > 0$$

cannot be improved: there exists a sequence (properly chosen linear combination of checkerboards on the macroelements) $\{\tilde{v}_h\}$ such that

$$|(\operatorname{div} \tau, \tilde{v}_h)| \leq C \|\tau\|_{L^2} \|\tilde{v}_h\|_{L^2}$$

Take $g_h = \tilde{v}_h / \|\tilde{v}_h\|_{L^2}$, then $g_h \rightharpoonup 0$ weakly in L^2 and $Tg_h \rightarrow 0$ strongly in L^2 .

Let's prove that $\|T_h g_h\|_{L^2}$ is bounded below away from zero (uniformly).

$$(\sigma_h, \tau) + (\operatorname{div} \tau, u_h) = 0 \quad \forall \tau \in \Sigma_h$$

$$(\operatorname{div} \sigma_h, v) = -(g_h, v) \quad \forall v \in U_h$$

$$(\sigma_h, \tau) + (\operatorname{div} \tau, u_h) = 0 \quad \forall \tau \in \Sigma_h$$

$$(\operatorname{div} \sigma_h, v) = -(g_h, v) \quad \forall v \in U_h$$

$$|(\operatorname{div} \sigma_h, u_h)| = |(g_h, u_h)| \leq \|u_h\|_{L^2}$$

$$(\sigma_h, \tau) + (\operatorname{div} \tau, u_h) = 0 \quad \forall \tau \in \Sigma_h$$

$$(\operatorname{div} \sigma_h, v) = -(g_h, v) \quad \forall v \in U_h$$

$$|(\operatorname{div} \sigma_h, u_h)| = |(g_h, u_h)| \leq \|u_h\|_{L^2}$$

$$|(\operatorname{div} \sigma_h, u_h)| = \|\sigma_h\|_{L^2}^2$$

$$(\sigma_h, \tau) + (\operatorname{div} \tau, u_h) = 0 \quad \forall \tau \in \Sigma_h$$

$$(\operatorname{div} \sigma_h, v) = -(g_h, v) \quad \forall v \in U_h$$

$$|(\operatorname{div} \sigma_h, u_h)| = |(g_h, u_h)| \leq \|u_h\|_{L^2}$$

$$|(\operatorname{div} \sigma_h, u_h)| = \|\sigma_h\|_{L^2}^2$$

$$\|g_h\|_{L^2} \|\sigma_h\|_{L^2} \geq \frac{1}{C} |(\operatorname{div} \sigma_h, g_h)|$$

$$(\sigma_h, \tau) + (\operatorname{div} \tau, u_h) = 0 \quad \forall \tau \in \Sigma_h$$

$$(\operatorname{div} \sigma_h, v) = -(g_h, v) \quad \forall v \in U_h$$

$$|(\operatorname{div} \sigma_h, u_h)| = |(g_h, u_h)| \leq \|u_h\|_{L^2}$$

$$|(\operatorname{div} \sigma_h, u_h)| = \|\sigma_h\|_{L^2}^2$$

$$\|g_h\|_{L^2} \|\sigma_h\|_{L^2} \geq \frac{1}{C} |(\operatorname{div} \sigma_h, g_h)| = \frac{1}{C} (g_h, g_h) = \frac{1}{C}$$

$$(\sigma_h, \tau) + (\operatorname{div} \tau, u_h) = 0 \quad \forall \tau \in \Sigma_h$$

$$(\operatorname{div} \sigma_h, v) = -(g_h, v) \quad \forall v \in U_h$$

$$|(\operatorname{div} \sigma_h, u_h)| = |(g_h, u_h)| \leq \|u_h\|_{L^2}$$

$$|(\operatorname{div} \sigma_h, u_h)| = \|\sigma_h\|_{L^2}^2$$

$$\|g_h\|_{L^2} \|\sigma_h\|_{L^2} \geq \frac{1}{C} |(\operatorname{div} \sigma_h, g_h)| = \frac{1}{C} (g_h, g_h) = \frac{1}{C}$$

$$\|u_h\|_{L^2} \geq \frac{1}{C^2}$$

<Back>

$\begin{pmatrix} 0 \\ g \end{pmatrix}$ case

$$U \subset H \simeq H' \subset U'$$

$\begin{pmatrix} 0 \\ g \end{pmatrix}$ case

$$U \subset H \simeq H' \subset U'$$

Weak approximability of U_H^0 with respect to $a(\cdot, \cdot)$.

$$b(\tau_h, u) \leq \rho(h) \|u\|_{U_H^0} \|\tau_h\|_a \quad \forall u \in U_H^0, \quad \forall \tau_h \in \ker_h(B)$$

$\begin{pmatrix} 0 \\ g \end{pmatrix}$ case

$$U \subset H \simeq H' \subset U'$$

Weak approximability of U_H^0 with respect to $a(\cdot, \cdot)$.

$$b(\tau_h, u) \leq \rho(h) \|u\|_{U_H^0} \|\tau_h\|_a \quad \forall u \in U_H^0, \quad \forall \tau_h \in \ker_h(B)$$

Strong approximability of U_H^0 .

For each $u \in U_H^0$ there exists $u^I \in U_h$ such that

$$\|u - u^I\| \leq \rho(h) \|u\|_{U_H^0}$$

$\begin{pmatrix} 0 \\ g \end{pmatrix}$ case

$$U \subset H \simeq H' \subset U'$$

Weak approximability of U_H^0 with respect to $a(\cdot, \cdot)$.

$$b(\tau_h, u) \leq \rho(h) \|u\|_{U_H^0} \|\tau_h\|_a \quad \forall u \in U_H^0, \quad \forall \tau_h \in \ker_h(B)$$

Strong approximability of U_H^0 .

For each $u \in U_H^0$ there exists $u^I \in U_h$ such that

$$\|u - u^I\| \leq \rho(h) \|u\|_{U_H^0}$$

+ Fortid

$\begin{pmatrix} \mathbf{f} \\ 0 \end{pmatrix}$ case

$$\Sigma \subset H \simeq H' \subset \Sigma'$$

$\begin{pmatrix} \mathbf{f} \\ 0 \end{pmatrix}$ case

$$\Sigma \subset H \simeq H' \subset \Sigma'$$

Weak approximability of U_0^H .

$$\sup_{\tau_h \in \ker_h(B)} \frac{b(\tau_h, u)}{\|\tau_h\|_\Sigma} \leq \rho(h) \|u\|_{U_0^H}$$

$\begin{pmatrix} \mathbf{f} \\ 0 \end{pmatrix}$ case

$$\Sigma \subset H \simeq H' \subset \Sigma'$$

Weak approximability of U_0^H .

$$\sup_{\tau_h \in \ker_h(B)} \frac{b(\tau_h, u)}{\|\tau_h\|_\Sigma} \leq \rho(h) \|u\|_{U_0^H}$$

Strong approximability of Σ_0^H .

For each $\sigma \in \Sigma_0^H$ there exists $\sigma^I \in \ker_h(B)$ such that

$$\|\sigma - \sigma^I\|_\Sigma \leq \rho(h) \|\sigma\|_{\Sigma_0^H}$$

$\begin{pmatrix} f \\ 0 \end{pmatrix}$ case

$$\Sigma \subset H \simeq H' \subset \Sigma'$$

Weak approximability of U_0^H .

$$\sup_{\tau_h \in \ker_h(B)} \frac{b(\tau_h, u)}{\|\tau_h\|_\Sigma} \leq \rho(h) \|u\|_{U_0^H}$$

Strong approximability of Σ_0^H .

For each $\sigma \in \Sigma_0^H$ there exists $\sigma^I \in \ker_h(B)$ such that

$$\|\sigma - \sigma^I\|_\Sigma \leq \rho(h) \|\sigma\|_{\Sigma_0^H}$$

+ DEK

<Back>