

Algebraic & Geometric Topology Volume 24 (2024)

Lattices, injective metrics and the $K(\pi, 1)$ conjecture

THOMAS HAETTEL





Lattices, injective metrics and the $K(\pi, 1)$ conjecture

THOMAS HAETTEL

Starting with a lattice with an action of \mathbb{Z} or \mathbb{R} , we build a Helly graph or an injective metric space. We deduce that the ℓ^{∞} orthoscheme complex of any bounded graded lattice is injective. We also prove a Cartan–Hadamard result for locally injective metric spaces. We apply this to show that any Garside group or any FC-type Artin group acts on an injective metric space and on a Helly graph. We also deduce that the natural piecewise ℓ^{∞} metric on any Euclidean building of type \tilde{A}_n extended, \tilde{B}_n , \tilde{C}_n or \tilde{D}_n is injective, and its thickening is a Helly graph.

Concerning Artin groups of Euclidean types \tilde{A}_n and \tilde{C}_n , we show that the natural piecewise ℓ^{∞} metric on the Deligne complex is injective, the thickening is a Helly graph, and it admits a convex bicombing. This gives a metric proof of the $K(\pi, 1)$ conjecture, as well as several other consequences usually known when the Deligne complex has a CAT(0) metric.

20E42, 05B35, 52A35, 06A12

Inti	Introduction	
1.	Review of posets, Artin groups and injective metric spaces	4013
2.	The thickening of a lattice	4022
3.	The affine version of a lattice	4025
4.	Application to Euclidean buildings and the Deligne complex of type \tilde{A}_n	4038
5.	The lattice of maximal parabolic subgroups of braid groups	4043
6.	The thickening of a semilattice	4047
7.	Application to Euclidean buildings and the Deligne complex of other Euclidean types	4051
8.	Bicombings on Deligne complexes in types \tilde{A}_n and \tilde{C}_n	4055
References		4057

Introduction

Injective metric spaces are geodesic metric spaces where every family of pairwise intersecting balls has a nonempty global intersection. The discrete counterpart of the injective metric space is the Helly graph. Its use in geometric group theory is recent and growing; see notably [Dress 1984; Lang 2013; Huang and Osajda 2021; Chalopin et al. 2020a; Haettel et al. 2023; Osajda and Valiunas 2024; Haettel 2022a]. Roughly speaking, CAT(0) spaces are typically locally Euclidean spaces, whereas injective metric

^{© 2024} MSP (Mathematical Sciences Publishers). Distributed under the Creative Commons Attribution License 4.0 (CC BY). Open Access made possible by subscribing institutions via Subscribe to Open.

Thomas Haettel

spaces are typically locally ℓ^{∞} metric spaces. Injective metric spaces display many nonpositive curvature features observed in CAT(0) spaces. We believe that Helly graphs and injective metric spaces may also prove to be more powerful than CAT(0) spaces for some purposes.

Firstly, it appears that many nonpositively curved groups (notably most CAT(0) groups) have a nice action on an injective metric space, so that the theory encompasses a vast class of groups: CAT(0) cubical groups, hyperbolic groups, relatively hyperbolic groups, uniform lattices in semisimple Lie groups over local fields, braid groups and more generally Garside groups, Artin groups of type FC, mapping class groups and more generally hierarchically hyperbolic groups. For instance, Helly graphs admit a simple combinatorial local characterization (see Theorem 1.12), which makes them potentially as powerful as CAT(0) cube complexes, whereas piecewise Euclidean CAT(0) complexes desperately lack a combinatorial characterization.

Secondly, we can still deduce many of the results that hold true for CAT(0) spaces and CAT(0) groups in the setting of injective spaces and injective groups (groups acting geometrically on injective spaces). For instance, injective metric spaces have a conical geodesic bicombing and are thus contractible, and every isometric bounded group action has a fixed point. Moreover, every injective group is semihyperbolic (see [Alonso and Bridson 1995]) and satisfies the Farrell–Jones conjecture (see [Kasprowski and Rüping 2017]). Sometimes we can even deduce stronger results than in the CAT(0) setting: as a sample result, note that any group acting properly and cocompactly on a Helly graph is biautomatic (see [Chalopin et al. 2020a]), whereas not all CAT(0) groups are biautomatic (see [Leary and Minasyan 2021]). Also note that infinite, finitely generated torsion groups do not act properly on uniformly locally finite Helly graphs (see [Haettel and Osajda 2021]), whereas the analogous statement is open for CAT(0) complexes.

In this article, we will pursue this philosophy. On one hand, we develop results useful to prove that some metric spaces are injective. On the other hand, we apply these results to Euclidean buildings and Deligne complexes of Euclidean Artin groups. We believe, however, that the scope of our results will not be limited to Artin groups of Euclidean type, and could concern much larger classes of groups.

We now present a very simple criterion (already appearing in a restricted form in [Haettel 2022a]) showing how to build a Helly graph or an injective metric space, starting with a lattice and an action of \mathbb{Z} or \mathbb{R} .

Theorem A (Theorem 2.1) Assume that *L* is a lattice such that each upper bounded subset of *L* has a join. Assume that there is an order-preserving increasing continuous action $(f_t)_{t \in H}$ of $H = \mathbb{Z}$ or \mathbb{R} on *L* such that,

for all
$$x, y \in L$$
, there exists $t \in H_+$ such that $f_{-t}(x) \leq y \leq f_t(x)$.

Let us define the following metric *d* on *L*:

for all $x, y \in L$, $d(x, y) = \inf\{t \in H_+ \mid f_{-t}(x) \le y \le f_t(x)\}.$

- If $H = \mathbb{Z}$, the metric space (L, d) is the vertex set of a Helly graph.
- If $H = \mathbb{R}$, the metric space (L, d) is injective.

Algebraic & Geometric Topology, Volume 24 (2024)

With Jingyin Huang, we used the same situation of a lattice with an appropriate action of \mathbb{Z} to define another natural graph which is weakly modular, and this applies to numerous examples (see [Haettel and Huang 2024]).

We applied this criterion in [Haettel 2022a] to prove that the thickening of a Bruhat–Tits Euclidean building of extended type \tilde{A}_n is a Helly graph. It turns out that the Bruhat–Tits restriction is unnecessary.

Theorem B (Theorems 4.4 and 7.4) The natural ℓ^{∞} metric on any Euclidean building of type \tilde{A}_n extended, \tilde{B}_n , \tilde{C}_n or \tilde{D}_n is injective. Furthermore, the thickening of its vertex set is a Helly graph.

Since Euclidean buildings are also endowed with CAT(0) metrics, we may wonder about the importance of this result. For one, it gives a lot of new examples to the theory of injective metric spaces and Helly graphs. Another consequence is another approach to Świątkowski's result [2006] that cocompact lattices in Euclidean buildings are biautomatic, in types \tilde{A}_n , \tilde{B}_n , \tilde{C}_n or \tilde{D}_n . The proof for types \tilde{B}_n , \tilde{C}_n and \tilde{D}_n relies on a generalization for graded semilattices; see Section 6.

Another immediate consequence of Theorem A is the following.

Theorem C (Corollary 2.2) The thickening of the Cayley graph of any Garside group with respect to its simple elements is a Helly graph.

Note that, in the case of a finite-type Garside group, this is due to Huang and Osajda [2021]. However, our proof is different, and does not rely on the deep local-to-global result for Helly graphs (see Theorem 1.12). Additionally, it also works in the case of a Garside group with infinite set of simple elements.

One application is the study of the orthoscheme complex of a bounded, graded lattice L. Note that simplices of the geometric realization |L| of L correspond to chains in L, so that one can endow each simplex with metric of the standard ℓ^{∞} orthosimplex associated to this order on vertices (see Section 1 for details). We endow the geometric realization |L| of L with a lattice structure, and use Theorem A to deduce the following.

Theorem D (Theorem 3.10) Let L denote a bounded, graded lattice. Then the orthoscheme complex |L| of L, with the piecewise ℓ^{∞} metric, is injective.

When L is a bounded, graded lattice, one may endow its orthoscheme complex with the piecewise Euclidean metric. Deciding whether it is a CAT(0) metric space is a very difficult question. It turns out that for orthoscheme complexes of posets, the CAT(0) property is more restrictive than the injective property; see Theorem 3.11.

One famous conjecturally CAT(0) example is the dual braid complex, defined by Brady and McCammond [2010]. The *n*-strand braid group B_n has a standard Garside structure, associated to the "half-turn" as a Garside element. The braid group also enjoys a dual presentation introduced by Birman, Ko and Lee [Birman et al. 1998], which corresponds to a dual Garside structure (see [Dehornoy and Paris 1999]). It is

Thomas Haettel

associated to the rotation of an n^{th} of a turn as a Garside element. Associated to this dual Garside structure, one may consider the geometric realization and endow it with the piecewise Euclidean orthoscheme metric (see Section 1). Brady and McCammond conjecture that this so-called dual braid complex is CAT(0) for all braid groups, but it has only been proved for $n \leq 7$ (see [Brady and McCammond 2010; Haettel et al. 2016; Jeong 2023]). However, an immediate consequence of Theorem D is the following.

Theorem E (Corollary 3.12) The Garside complex of any Garside group, endowed with the piecewise ℓ^{∞} orthoscheme metric, is injective.

We also obtain another proof of a result by Huang and Osajda stating that FC-type Artin groups are Helly [Huang and Osajda 2021, Theorem 5.8]. We also provide explicit Helly and injective models (see Theorem 7.6 for the precise statement).

Theorem F (Huang–Osajda, Theorem 7.6) Let A denote an Artin group of type FC. Then a natural simplicial complex X with vertex set A, with the ℓ^{∞} metric, is injective. Moreover, a thickening of X is a Helly graph.

As a particular case of Theorem E, the dual braid complex, endowed with the piecewise ℓ^{∞} orthoscheme metric, is injective for all braid groups. This also holds, more generally, for every spherical-type Artin group with some Garside structure. Note that Theorem D proves local injectivity, and one also needs a Cartan–Hadamard result for injective metric spaces in order to conclude. We therefore rely on the local-to-global result for Helly graphs to prove the following generalization of [Miesch 2017] in the nonproper setting (see Section 1 for the definition of semiuniformly locally injective). This result is clearly of independent interest.

Theorem G (Cartan–Hadamard for injective metric spaces, Theorem 1.14) Let X denote a complete, simply connected, semiuniformly locally injective metric space. Then X is injective.

Another very promising family of examples is Deligne complexes of Artin groups (see Section 1 for definitions). Note that buildings and Deligne complexes of Artin groups are closely related: in fact in his original article Deligne [1972] called "buildings for generalized braid groups" the complexes later called Deligne complexes. However, to the best of our knowledge, the close relationship between Euclidean buildings and Deligne complexes of Euclidean-type Artin groups has not yet been exploited in the literature. Notably, the automorphism groups of Euclidean buildings do not possess a Garside structure, and the Deligne complexes do not have an apartment system as rich as in the building case. However, one common feature is that they locally look like a lattice, which is the key combinatorial property we are using in this article.

Associated to every Coxeter graph Γ with vertex set *S*, we may define the Coxeter group $W(\Gamma)$, the Artin group $A(\Gamma)$ and the hyperplane complement $M(\Gamma)$ (see Section 1). The Coxeter group $W(\Gamma)$ acts naturally on $M(\Gamma)$, and $A(\Gamma)$ is the fundamental group of the quotient $W(\Gamma) \setminus M(\Gamma)$. One very natural question is to decide whether it is a classifying space. This is the statement of the following conjecture.

Algebraic & Geometric Topology, Volume 24 (2024)

Conjecture H (the $K(\pi, 1)$ conjecture) The hyperplane complement $M(\Gamma)$ is aspherical.

The $K(\pi, 1)$ conjecture has been proved for spherical-type Artin groups by Deligne [1972], and for Euclidean-type Artin groups by Paolini and Salvetti [2021] very recently, even for the type \tilde{D}_n . Another approach, more closely related to the metric approach of this article, was used by Charney and Davis [1995b] to prove the $K(\pi, 1)$ conjecture for Artin groups of type FC or of 2–dimensional type. Their proof relies on the use of a simplicial complex, called the Deligne complex $\Delta(\Gamma)$, which is the geometric realization of the poset of cosets of spherical-type parabolic subgroups (see Section 1). This complex (in this form) was defined in [Charney and Davis 1995b], where they proved that $\Delta(\Gamma)$ is homotopy equivalent to the universal cover of $W(\Gamma) \setminus M(\Gamma)$. In other words, the $K(\pi, 1)$ conjecture is equivalent to the contractibility of the Deligne complex $\Delta(\Gamma)$.

Charney and Davis's method for proving that the Deligne complex is contractible is to endow it with a CAT(0) metric. This works for Artin groups of type FC or of 2-dimensional type. However, in the general case, the key question is to decide whether the Deligne complex for the braid group is CAT(0) with Moussong's metric. It is only known for the braid group up to four strands only (see [Charney 2004]). However, we will see that the natural piecewise ℓ^{∞} metric is injective for all braid groups, up to taking the product with \mathbb{R} .

Theorem I (Theorem 4.4) Let Δ denote the Deligne complex of the Artin group of Euclidean type \tilde{A}_n . Then the natural piecewise ℓ^{∞} length metric on $\Delta \times \mathbb{R}$ is injective. Moreover, the thickening of $\Delta \times \mathbb{R}$ is a Helly graph.

The proof consists in applying Theorem D to prove that the Deligne complex is locally injective. One key combinatorial property is that the Deligne complex in type A_n is essentially a lattice; see Section 5. The proof of the lattice property, through the cut-curve lattice, is due to Crisp and McCammond, copied here with their permission.

Note that one may wonder whether it is necessary to consider the direct product with \mathbb{R} . In fact, Hoda [2023] proved that the Euclidean Coxeter group of type \tilde{A}_n is not Helly for $n \ge 2$, even though its direct product with \mathbb{Z} is. We made a similar distinction for automorphism groups of Euclidean buildings of type \tilde{A}_n in [Haettel 2022a]. We therefore strongly believe that there is no injective metric on the Deligne complex of type \tilde{A}_n itself which is invariant under the Artin group. However, we will see in Theorem M below that there is a convex bicombing on the Deligne complex itself.

In order to deal with the Euclidean type \tilde{C}_n , we first prove a generalization of Theorem D for graded semilattices; see Section 6.

Theorem J (Theorem 7.4) Let Δ denote the Deligne complex of the Artin group of Euclidean type \tilde{C}_n . Then the natural piecewise ℓ^{∞} length metric on Δ is injective. Moreover, the thickening of Δ is a Helly graph.

An immediate consequence is another proof of the $K(\pi, 1)$ conjecture in Euclidean types \tilde{A}_n and \tilde{C}_n , originally due to Okonek [1979]. The novelty is that it is the first metric proof. Moreover, in Charney and Davis's approach to the $K(\pi, 1)$ conjecture by showing that Moussong's metric is CAT(0), the main difficulty is to prove that it is locally CAT(0) for the braid groups. It is precisely this statement that we are able to prove in the injective setting.

Corollary K (Okonek) The Deligne complex Δ of Euclidean type \tilde{A}_n or \tilde{C}_n is contractible. In particular, the $K(\pi, 1)$ conjecture holds in these cases.

Moreover, several results have been proved, relying on the assumption that one may endow the Deligne complex with a piecewise Euclidean CAT(0) metric. Crisp [2000] studied the fixed-point subgroup under a symmetry group of the Artin system. Godelle [2007] studied the centralizer and normalizer of standard parabolic subgroups. Morris-Wright [2021] studied the intersections of parabolic subgroups. It turns out almost all the arguments merely used the existence of an equivariant geodesic bicombing on the Deligne complex (see the end of Section 3 for a definition of a bicombing). Moreover, Descombes and Lang [2015; 2016] justified the importance of convex geodesic bicombings for themselves. We therefore state the following conjecture, which may be seen as a metric strategy for the proof of the $K(\pi, 1)$ conjecture.

Conjecture L The Deligne complex of any Artin group *A* has an *A*–invariant metric that admits a convex, consistent, reversible geodesic bicombing.

We are able to prove this conjecture in spherical types A_n , B_n and Euclidean types \tilde{A}_n , \tilde{C}_n , and we believe that our result represents a major step towards the general case.

Theorem M (Theorem 8.1) Let Δ denote the Deligne complex of the Artin group of spherical type A_n or B_n or Euclidean type \tilde{A}_n or \tilde{C}_n . There exists a metric on Δ , invariant under the Artin group, which admits a convex, consistent, reversible geodesic bicombing.

If an Artin group satisfies Conjecture L, then the $K(\pi, 1)$ conjecture follows. Moreover, we may also list consequences of [Crisp 2000; Godelle 2007; Morris-Wright 2021; Cumplido et al. 2019] that rely on the assumption that the Deligne complex has a CAT(0) metric. However, most of the arguments only use the geodesic bicombing. The following results were only essentially known for Artin groups of type FC or of 2–dimensional type. Note that concerning Artin groups of type FC, only intersections of spherical-type parabolic subgroups are known to be parabolic (see [Morris-Wright 2021]). See also [Möller et al. 2024].

Corollary N (Corollaries 8.3, 8.4 and 8.5) Let A denote the Artin group of Euclidean type \tilde{A}_n or \tilde{C}_n .

- The intersection of any parabolic subgroups of A is a parabolic subgroup.
- *A* satisfies Properties (★), (★★) and (★★★) from [Godelle 2007], notably: for any subset *X* ⊂ *S*, we have

 $\operatorname{Com}_{A}(A_{X}) = N_{A}(A_{X}) = A_{X} \cdot QZ_{A}(X),$

where the quasicentralizer of X is $QZ_A(X) = \{g \in A \mid g \cdot X = X\}.$

Algebraic & Geometric Topology, Volume 24 (2024)

• For any group G of symmetries of the Artin system, the fixed-point subgroup A^G is isomorphic to an Artin group.

In view of Conjecture L, looking for other consequences of the existence of a convex bicombing on the Deligne complex may prove to be fruitful.

Structure of the article

In Section 1, we review basic definitions of posets, lattices, Artin groups, Deligne complexes, injective metric spaces and Helly graphs. We also prove the Cartan–Hadamard theorem for injective metric spaces. In Section 2, we prove the central simple criterion showing how to produce an injective metric space or a Helly graph starting from a lattice with an action of \mathbb{Z} or \mathbb{R} . In Section 3, we apply this criterion to prove that the orthoscheme complex of a bounded, graded lattice is injective. In Section 4, we use this to prove that, for Euclidean buildings and the Deligne complex in Euclidean type \tilde{A}_n , the natural piecewise ℓ^{∞} metric is injective. In Section 6, we show how to adapt the criterion to a mere semilattice with some extra property. We then apply it in Section 7 to prove that, for Euclidean buildings and the Deligne complex in Euclidean type \tilde{C}_n , the natural piecewise ℓ^{∞} metric is injective. Finally, in Section 8, we use the convex bicombing on the Deligne complexes to deduce many corollaries about parabolic subgroups of Artin groups.

Acknowledgments

We would like to thank very warmly Jingyin Huang, Jon McCammond, Damian Osajda and Luis Paris for very interesting discussions. We are indebted to Jon McCammond and John Crisp for their authorization to copy their work about the lattice of cut-curves. We also would like to thank the organizers of the ICMS Edinburgh 2021 meeting *Perspectives on Artin groups*, where we had fruitful discussions. We also would like to thank Jérémie Chalopin, Victor Chepoi, María Cumplido, Anthony Genevois, Hiroshi Hirai, Alexandre Martin and Bert Wiest for various discussions. We also thank an anonymous referee for many helpful comments.

We acknowledge support from French project ANR-16-CE40-0022-01 AGIRA.

1 Review of posets, Artin groups and injective metric spaces

1.1 Posets and lattices

We recall briefly basic definitions related to posets and lattices.

Definition 1.1 (poset) Let *L* denote a poset. A *chain* of *L* is a totally ordered nonempty subset of *L*. A *maximal chain* is a chain that is maximal with respect to inclusion. A finite chain of n + 1 elements $x_0 < x_1 < \cdots < x_n$ is called of *length n*. The poset *L* is called *bounded below* (resp. above) if it has a global minimum denoted by 0 (resp. global maximum denoted by 1). The poset *L* is called *bounded* if it is both bounded above and below. The poset *L* has *rank n* if it is bounded and all maximal chains have length *n*.

Definition 1.2 (interval) Given two elements $x \le y$ in a poset L, we define the *interval*

$$I(x, y) = \{ z \in L \mid x \le z \le y \}.$$

In the case of possible confusion, we may also denote the interval by $I_L(x, y)$ to emphasize that we are considering the interval in the poset L. The poset L is called *graded* if every interval of L has a rank. Let x denote an element in a graded poset L that is bounded below; the *rank* of x is the rank of the interval I(0, x).

Definition 1.3 (lattice) Given elements x, y in a poset L, if there exists a unique maximal lower bound to $\{x, y\}$, it is called the *meet* of x and y and denoted by $x \land y$. Similarly, if there exists a unique minimal upper bound to $\{x, y\}$, it is called the *join* of x and y and denoted by $x \lor y$. If any two elements of L have a meet (resp. a join), the poset L is called a *meet-semilattice* (resp. join-semilattice). If any two elements of L have a meet and a join, the poset L is called a *lattice*.

Examples • The Boolean lattice *L* of rank *n* is the poset of subsets of $E = \{1, ..., n\}$, partially ordered by inclusion. The join of $A, B \in L$ is $A \cup B$ and their meet is $A \cap B$.

• Consider a CAT(0) cube complex X, with a base vertex v_0 . Order the set V of vertices of X by declaring that $v \leq w$ if some combinatorial geodesic from v_0 to w passes through v. Then V is a graded meet-semilattice, with minimum v_0 . The meet of two vertices $v, w \in V$ is the median of v_0, v, w .

• The partition lattice *L* of $E = \{1, ..., n\}$ is the poset of partitions of *E*, partially ordered by declaring that $A \leq B$ if every element of *A* is contained in an element of *B*. This lattice has rank n-1, its minimum is the partition $\{\{1\}, \{2\}, ..., \{n\}\}$ into singletons and its maximum is the partition $\{E\}$ into one element.

We will now describe a very simple criterion due to [Brady and McCammond 2010] to decide when a bounded graded poset is a lattice.

Definition 1.4 (bowtie) In a poset *L*, a *bowtie* consists of four distinct elements *a*, *b*, *c*, *d* such that *a*, *c* are minimal upper bounds of *b*, *d*, and *b*, *d* are maximal lower bounds of *a*, *c*.

Proposition 1.5 [Brady and McCammond 2010, Proposition 1.5] Let *L* denote a bounded graded poset. Then *L* is a lattice if and only if *L* does not contain a bowtie.

1.2 Coxeter groups, Artin groups and Deligne complexes

We recall the definitions of Coxeter groups, Artin groups, and their associated Deligne complexes.

For every finite simple graph Γ with vertex set *S* and with edges labeled by some integer in $\{2, 3, ...\}$, one associates the Coxeter group $W(\Gamma)$ with the following presentation:

 $W(\Gamma) = \langle S \mid \text{for all } \{s, t\} \in \Gamma^{(1)}, \text{ for all } s \in S, \ s^2 = 1, \ [s, t]_m = [t, s]_m \text{ if the edge } \{s, t\} \text{ is labeled } m \rangle,$

where $[s, t]_m$ denotes the word ststs... of length m.

 ${\cal A} {\it lgebraic} \,\, {\it \&Geometric} \,\, {\it Topology, Volume} \,\, 24 \, (2024)$

	spherical type		Euclidean type
$A_n, n \ge 2$	••	••	\tilde{A}_n • • • • • • • • • • • • • • • • • • •
$B_n, n \ge 2$	••	•••	\widetilde{B}_n $\overset{\bullet}{\longrightarrow}$ $\overset{\bullet}{\longrightarrow}$ $\overset{\bullet}{\longrightarrow}$ $\overset{\bullet}{\longrightarrow}$ $\overset{\bullet}{\longrightarrow}$ $\overset{\bullet}{\longrightarrow}$
$C_n = B_n, n \ge 2$	•••	••_4_•	$\widetilde{C}_n \bullet \overset{4}{-} \bullet \cdots \bullet \overset{-}{-} \bullet \overset{4}{-} \bullet$
$D_n, n \ge 4$	••	•—•<	\tilde{D}_n \rightarrow \cdots \cdot \cdot



The associated Artin group $A(\Gamma)$ is defined by a similar presentation:

 $A(\Gamma) = \langle S \mid \text{for all } \{s, t\} \in \Gamma^{(1)}, \ [s, t]_m = [t, s]_m \text{ if the edge } \{s, t\} \text{ is labeled } m \rangle.$

The groups $A(\Gamma)$ are also called Artin–Tits groups, since they were defined by Tits [1966].

Note that only the relations $s^2 = 1$ have been removed, so that there is a natural surjective morphism from $A(\Gamma)$ to $W(\Gamma)$. Also note that if m = 2, then s and t commute, and if m = 3, then s and t satisfy the classical braid relation sts = tst.

The knowledge of general Artin groups is extremely limited (see notably [McCammond 2017; Charney 2008; Godelle and Paris 2012]). In particular, we do not know whether the word problem is solvable in general Artin groups, nor whether they are torsion-free.

Most results about Artin–Tits groups concern particular classes. The Artin group $A(\Gamma)$ is called:

- of *spherical type* if its associated Coxeter group $W(\Gamma)$ is finite, it may be realized as a reflection group of a sphere.
- of *Euclidean type* if its associated Coxeter group $W(\Gamma)$ may be realized as a reflection group of a Euclidean space.
- of *FC type* if for any complete subset $T \subset S$ the parabolic subgroup $A_T = \langle T \rangle$ is spherical.

We recall in Table 1 the classification of the four infinite families of spherical and Euclidean irreducible diagrams; see [Bourbaki 2002] for the full classification. We only present those because we will only consider these types in this article. Note that we use in this table the convention of Dynkin diagrams: vertices that are not joined by an edge commute, and we drop the label 3 from edges.

Artin groups are closely related to hyperplane complements, which can be presented in a simple way in spherical and Euclidean types. Fix a Coxeter group $W = W(\Gamma)$ of spherical type or Euclidean type acting by reflections on a sphere \mathbb{S}^{n-1} or a Euclidean space \mathbb{R}^n . In the case W is of spherical type, consider the

associated linear action on \mathbb{R}^n . A conjugate of an element of the standard generating set *S* is called a *reflection*. Let \mathcal{R} denote the set of reflections in *W*. Consider the family of affine hyperplanes of \mathbb{R}^n

$$\mathcal{H} = \{ H_r \mid r \in \mathcal{R} \},\$$

where $H_r \subset \mathbb{R}^n$ denotes the fixed-point set of the reflection *r*.

The complement of the complexified hyperplane arrangement is

$$M(\Gamma) = \mathbb{C}^n \setminus \bigcup_{r \in \mathcal{R}} (\mathbb{C} \otimes H_r).$$

Note that W acts naturally on M, and we have the following (see [van der Lek 1983]):

$$\pi_1(W(\Gamma)\backslash M(\Gamma)) \simeq A(\Gamma).$$

So the Artin group $A(\Gamma)$ appears as the fundamental group of (a quotient of) the complement of a complexified hyperplane arrangement. One very natural question is to decide whether it is a classifying space. This is the statement of the following conjecture.

Conjecture (the $K(\pi, 1)$ conjecture) The space $M(\Gamma)$ is aspherical.

This conjecture was proved for spherical-type Artin groups by Deligne [1972], and for Euclidean-type Artin groups by Paolini and Salvetti [2021] very recently, even for the type \tilde{D}_n . Another approach, more closely related to the content of this article, was used by Charney and Davis [1995a] to prove the $K(\pi, 1)$ conjecture for Artin groups of type FC or of 2-dimensional type. Their proof relies on the use of a simplicial complex, called the Deligne complex, and they endow it with a particular metric to show that it is contractible.

We will now recall the definition of the Deligne complex of an Artin group $A = A(\Gamma)$. A standard parabolic subgroup of A is the subgroup $A_T = \langle T \rangle$ generated by a subset T of S, the standard generating set of A. A parabolic subgroup denotes any conjugate of a standard parabolic subgroup. Let us define

$$S_f = \{T \subset S \mid W_T \text{ is finite}\}.$$

The *Deligne complex* $\Delta = \Delta(\Gamma)$ is the order complex of the set of cosets of parabolic subgroups

$$\{gA_T \mid g \in A, T \in \mathcal{S}_f\},\$$

where the partial order is given by the inclusion $gA_T \subset g'A_{T'}$ of cosets.

One key property of the Deligne complex is that it has the same homotopy type as the universal cover of the hyperplane complement:

Theorem 1.6 [Charney and Davis 1995b, Theorem 1.5.1] The Deligne complex $\Delta(\Gamma)$ is homotopy equivalent to the universal cover of the quotient of the hyperplane complement $W(\Gamma) \setminus M(\Gamma)$.

In particular, the $K(\pi, 1)$ conjecture amounts to proving that the Deligne complex is contractible.

Algebraic & Geometric Topology, Volume 24 (2024)

1.3 Injective metrics and Helly graphs

We briefly recall basic definitions of injective metric spaces and Helly graphs. We will also state the local-to-global result for Helly graphs and deduce the analogous Cartan–Hadamard result for injective metric spaces.

A geodesic metric space is called *injective* (one may also say *hyperconvex*, or *absolute* 1–*Lipschitz retract*) if any family of pairwise intersecting closed balls has a nonempty global intersection, the so-called *Helly property*. We refer the reader to [Lang 2013] for a presentation of injective metric spaces.

Examples • The normed vector space $(\mathbb{R}^n, \ell^{\infty})$ is injective for all $n \ge 1$. In fact, it is up to isometry the only injective norm on \mathbb{R}^n (see [Nachbin 1950]).

- Any tree is injective.
- Any finite-dimensional CAT(0) cube complex, endowed with the standard piecewise ℓ^{∞} metric, is injective (see [Miesch 2014] and [Bowditch 2020]).

A connected graph X is called *Helly* if any family of pairwise intersecting combinatorial balls of X has a nonempty global intersection, in other words the combinatorial balls satisfy the Helly property. We refer the reader to [Chalopin et al. 2020a] for a presentation of Helly graphs and Helly groups. Many examples of Helly graphs come from thickening of complexes, which we define now (see [Chalopin et al. 2020a; Huang and Osajda 2021]).

Definition 1.7 (thickening) Let X denote a cell complex. The *thickening* of X (with respect to the cell structure) is the graph with vertex set $X^{(0)}$, with an edge between two vertices if and only if they are contained in a common cell of X.

Examples • For each $n \ge 1$, the graph with vertex set \mathbb{Z}^n , with an edge between v, w if $d_{\infty}(v, w) = 1$, is a Helly graph.

- Any simplicial tree is a Helly graph.
- The thickening of the vertex set of a CAT(0) cube complex is a Helly graph.

In order to endow a simplicial complex with a potentially injective (or CAT(0)) metric, it is natural to ask for metric simplices which may tile the Euclidean space \mathbb{R}^n . One choice is to consider the barycentric subdivision of the standard cubical tiling of \mathbb{R}^n , whose simplices are orthosimplices, which we now formally define.

Definition 1.8 (orthosimplex) The *standard orthosimplex* of dimension *n* is the simplex of \mathbb{R}^n with vertices $(0, \ldots, 0), (1, 0, \ldots, 0), \ldots, (1, 1, \ldots, 1)$ (see Figure 1). One may endow the simplex with the standard ℓ^p metric on \mathbb{R}^n for any $p \in [1, \infty]$. Throughout this article (except in Theorem 3.11), we will



Figure 1: The standard 3-dimensional orthosimplex.

only consider the ℓ^{∞} metric, called the ℓ^{∞} orthosimplex. Note that any *n*-simplex with a total order on its vertices $v_0 < v_1 < \cdots < v_n$ may be identified uniquely with the ℓ^{∞} orthosimplex of dimension *n*, where each v_i is identified with the vertex $(1, \ldots, 1, 0, \ldots, 0)$ with *i* ones and n - i zeros. Also note that reversing the total order on the vertices gives rise to an isometry of the orthosimplex.

Definition 1.9 (simplicial complex with ordered simplices and maximal edges) We say X is a simplicial complex X with ordered simplices if each simplex of X has a total order on its vertex set, which is consistent with respect to inclusions of simplices of X. Moreover, we say that X has maximal edges if, given any simplex σ of X, there exist adjacent vertices a, b in X such that $\sigma \cup \{a, b\}$ is a simplex of X, and such that any vertex c of X adjacent to both a and b satisfies $a \le c \le b$. Such an edge $\{a, b\}$ is then called a maximal edge of X.

Definition 1.10 (orthoscheme complex) Let X denote a finite-dimensional simplicial complex with ordered simplices. Then one may endow each simplex of X with the associated ℓ^{∞} orthoscheme metric. Then the geometric realization of |X|, endowed with the length metric associated with the ℓ^{∞} orthoscheme metric on each simplex, is called the ℓ^{∞} orthoscheme complex of X.

As a particular case, if L is a poset with a bound on the length of its chains, then the geometric realization of |L| satisfies the assumptions, so we may talk about the ℓ^{∞} orthoscheme complex of L. The following is an immediate adaptation of [Bridson and Haefliger 1999, Theorem 7.13].

Theorem 1.11 Let X denote a finite-dimensional simplicial complex with ordered simplices. Then its ℓ^{∞} orthoscheme complex is a complete length space.

Proof Note that since X has dimension n, the ℓ^{∞} orthoscheme complex of X has finitely many isometry types of cells: the standard orthoscheme k-simplices, where $k \leq n$. The proof of [Bridson and Haefliger 1999, Theorem 7.13] adapts without change to this situation.

One key property in the study of Helly graphs is the following local-to-global statement (see [Chalopin et al. 2020b]). Recall that a *clique* of a graph is a complete subgraph. A graph is called *clique-Helly* if its

family of maximal cliques satisfies the Helly property (ie any family of pairwise intersecting maximal cliques has nonempty intersection). The triangle complex of a simplicial graph X is the simplicial 2-complex whose 1-skeleton is X, and whose 2-simplices correspond to triangles in X.

Theorem 1.12 [Chalopin et al. 2020b, Theorem 3.5] Let *X* denote a graph. Then *X* is Helly if and only if *X* is clique-Helly and its triangle complex is simply connected.

In order to transfer this local-to-global property to injective metric spaces, we will need the following technical lemma. Say that a metric space is ε -coarsely injective (for some $\varepsilon \ge 0$) if, for any families $(x_i)_{i \in I}$ in X and $(r_i)_{i \in I}$ in \mathbb{R}_+ such that, for all $i, j \in I$, $d(x_i, x_j) \le r_i + r_j$, we have $\bigcap_{i \in I} B_X(x_i, r_i + \varepsilon) \ne \emptyset$. Note that when $\varepsilon = 0$, we recover the definition of an injective metric space.

Lemma 1.13 Let X denote a complete metric space that is ε -coarsely injective for every $\varepsilon > 0$. Then X is injective.

Proof We will prove that X is injective: consider a family $(B_X(x_i, r_i))_{i \in I}$ of pairwise intersecting balls in X. We know that, for all $i, j \in I$, $d(x_i, x_j) \leq r_i + r_j$. For any $\varepsilon > 0$, let us define $A_{\varepsilon} = \bigcap_{i \in I} B_X(x_i, r_i + \varepsilon)$, which is nonempty by the assumption of ε -coarse injectivity.

Fix $0 < \varepsilon \leq \varepsilon'$. We will prove that the Hausdorff distance between A_{ε} and $A_{\varepsilon'}$ is at most $\varepsilon + \varepsilon'$. Note that $A_{\varepsilon} \subset A_{\varepsilon'}$. Fix $x_0 \in A_{\varepsilon'}$. We will prove that $d(x_0, A_{\varepsilon}) \leq \varepsilon + \varepsilon'$. Let $I_0 = I \sqcup \{0\}$, and let $r_0 = \varepsilon'$. Consider the families $(x_i)_{i \in I_0}$ in X and $(r_i)_{i \in I_0}$ in \mathbb{R}_+ . For each $i, j \in I_0$, we know that $d(x_i, x_j) \leq r_i + r_j$: indeed, for any $i \in I$, we have $x_0 \in B_X(x_i, r_i + \varepsilon')$. By ε -coarse injectivity, we deduce that the intersection $\bigcap_{i \in I_0} B_X(x_i, r_i + \varepsilon)$ is not empty. In particular, the ball $B_X(x_0, \varepsilon + \varepsilon')$ intersects $A_{\varepsilon} = \bigcap_{i \in I} B_X(x_i, r_i + \varepsilon)$. This implies that $d(x_0, A_{\varepsilon}) \leq \varepsilon + \varepsilon'$. So we have proved that the Hausdorff distance between A_{ε} and $A_{\varepsilon'}$ is at most $\varepsilon + \varepsilon'$.

For each $n \in \mathbb{N}$, consider by induction $x_n \in A_{2^{-n}}$ such that, for all $n \ge 0$,

$$d_X(x_{n+1}, x_n) \leq 2^{-n} + 2^{-(n+1)} \leq 2^{-n+1}$$

For each $0 \le n \le m$, we have $d_X(x_n, x_m) \le 2^{-n+2}$; hence the sequence $(x_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in X. Since X is complete, it has a limit $y \in X$. For each $n \in \mathbb{N}$, we have $y \in A_{2^{-n}}$, so for each $i \in I$ we have $d_X(y, x_i) \le r_i + 2^{-n}$. We deduce that, for each $i \in I$, we have $d_X(y, x_i) \le r_i$. In other words, y belongs to the intersection $\bigcap_{i \in I} B(x_i, r_i)$: we have proved that X is injective.

Say that a metric space is *uniformly locally injective* if there exists $\varepsilon > 0$ such that each ball of radius ε is injective. Say that a metric space is *semiuniformly locally injective* if there exists $\varepsilon > 0$ such that each ball of radius ε is uniformly locally injective. For instance, any locally compact, locally injective metric space is semiuniformly locally injective.

As a concrete example, if X denotes the injective hull of the hyperbolic plane $\mathbb{H}^2_{\mathbb{R}}$ and Γ is a nonuniform lattice in PGL(2, \mathbb{R}) = Isom($\mathbb{H}^2_{\mathbb{R}}$), then the quotient $\Gamma \setminus X$ is semiuniformly locally injective, but not uniformly locally injective. There are similar examples in higher rank; see [Haettel 2022a].

There are also nonlocally compact examples: let X denote a metric simplicial graph such that the systole at each vertex of X is bounded below. Then X is locally uniformly locally injective. However, if the systole of X is 0, then X is not uniformly locally injective.

We now prove a Cartan–Hadamard theorem for such injective metric spaces, relying on the local-to-global property for Helly graphs, Theorem 1.12. Note that this statement generalizes [Miesch 2017, Theorem 1.2] without the local compactness assumption.

Theorem 1.14 (Cartan–Hadamard for injective metric spaces) Let X denote a complete, simply connected, semiuniformly locally injective metric space. Then X is injective.

Proof We will first prove the statement when X is uniformly locally injective. Fix $\varepsilon > 0$ small enough such that balls in X of radius at most 2ε are injective. Consider the graph Γ_{ε} with vertex set X and with an edge between $x, y \in X$ if $d(x, y) \leq \varepsilon$. Since X is geodesic, Γ_{ε} is a connected graph. Also note that, for any $x \in X$ and $n \in \mathbb{N}$, we have

$$B_{\Gamma_{\varepsilon}}(x,n) = B_X(x,n\varepsilon).$$

We will prove that, for each $\varepsilon > 0$, the graph Γ_{ε} is a Helly graph by applying Theorem 1.12.

We first prove that the family of combinatorial 1-balls in Γ_{ε} satisfies the Helly property. Fix a family of vertices $(x_i)_{i \in I}$ of Γ_{ε} such that, for all $i, j \in I$, $d_{\Gamma_{\varepsilon}}(x_i, x_j) \leq 2$. We want to prove that these balls intersect in Γ_{ε} .

The family of metric balls $(B_X(x_i, \varepsilon))_{i \in I}$ in X pairwise intersects: since such balls have the Helly property by the assumption on X, we deduce that there exists $y \in X$ such that, for all $i \in I$, $d_X(x_i, y) \leq \varepsilon$. In other words, the vertex $y \in \Gamma_{\varepsilon}$ lies in the intersection of all combinatorial 1-balls $(B_{\Gamma_{\varepsilon}}(x_i, 1))_{i \in I}$.

We now deduce that Γ_{ε} is clique-Helly: fix a family of pairwise intersecting maximal cliques $(\sigma_i)_{i \in I}$ of Γ_{ε} . Then the family of combinatorial 1-balls centered at each vertex of each clique σ_i , for $i \in I$, pairwise intersects: according to the previous paragraph, we deduce that there exists a vertex $y \in \Gamma_{\varepsilon}$ adjacent to each vertex of each clique σ_i for $i \in I$. Since each clique σ_i is maximal, we deduce that y belongs to the intersection of all σ_i for $i \in I$. The graph Γ_{ε} is clique-Helly.

We now prove that the triangle complex of Γ_{ε} is simply connected. Fix a combinatorial loop ℓ in Γ_{ε} . Since X is simply connected, there exists a disk D in X bounding ℓ . Consider a triangulation T of D such that triangles have diameter for d_X at most ε . Then the vertex set of each triangle of T is a clique in Γ_{ε} ; therefore ℓ is null-homotopic in the triangle complex of Γ_{ε} . So the triangle complex of Γ_{ε} is simply connected.

The graph Γ_{ε} is clique-Helly and has a simply connected triangle complex, so according to Theorem 1.12, we deduce that Γ_{ε} is a Helly graph.

Note that, for any $\varepsilon > 0$, we have $d_X \leq \varepsilon d_{\Gamma_{\varepsilon}} \leq d_X + \varepsilon$, and balls for the metric $d_{\Gamma_{\varepsilon}}$ with integral radius satisfy the Helly property.

We will show that the metric space X is ε -coarsely injective: consider families $(x_i)_{i \in I}$ in X and $(r_i)_{i \in I}$ in \mathbb{R}_+ such that, for all $i, j \in I$, $d_X(x_i, x_j) \leq r_i + r_j$. For each $i \in I$, let $n_i \in \mathbb{N}$ such that $n_i \varepsilon \leq r_i < (n_i + 1)\varepsilon$. For each $i, j \in I$, since the balls $B_X(x_i, r_i)$ and $B_X(x_j, r_j)$ intersect, we deduce that $d_{\Gamma_{\varepsilon}}(x_i, x_j) \leq n_i + n_j + 2$. So, in the Helly graph Γ_{ε} , the balls $(B_{\Gamma_{\varepsilon}}(x_i, n_i + 1))_{i \in I}$ pairwise intersect: we deduce that there exists $y \in \Gamma_{\varepsilon}$ such that, for all $i \in I$, $d_{\Gamma_{\varepsilon}}(y, x_i) \leq n_i + 1$. In particular, for any $i \in I$, we have $d_X(y, x_i) \leq (n_i + 1)\varepsilon \leq r_i + \varepsilon$. Hence y belongs to each ball $B_X(x_i, r_i + \varepsilon)$, for $i \in I$: this proves that X is ε -coarsely injective.

Since this holds for any small enough $\varepsilon > 0$, according to Lemma 1.13, we conclude that X is injective.

We now turn to the general case, when X is only semiuniformly locally injective: there exists $\varepsilon > 0$ such that each ball of radius ε in X is uniformly locally injective. According to the uniformly locally injective case, we deduce that each ball of radius ε is injective: this means that X is uniformly locally injective. According to the uniformly locally injective case again, we deduce that X is injective.

We now see that, under a mild assumption on a simplicial complex with ordered simplices, saying that the ℓ^{∞} orthoscheme realization is injective is equivalent to saying that the thickening of the 1–skeleton is Helly. Note that, if we refer to the definition of thickening as in Definition 1.7, the corresponding cell structure is not the simplicial one, but a coarser cell structure whose cells correspond to intervals.

Theorem 1.15 Let X denote a finite-dimensional simplicial complex with ordered simplices, and with maximal edges. Let Γ denote the graph with vertex set $X^{(0)}$, and with an edge between $x, y \in X^{(0)}$ if there exist $a, b \in X^{(0)}$ and ordered triangles $a \leq x \leq b$ and $a \leq y \leq b$ in X. If the ℓ^{∞} orthoscheme complex of X is injective, then the thickening Γ of X is a Helly graph.

Proof We see that maximal cliques in Γ correspond to intervals

 $I_{ab} = \{x \in X^{(0)} \mid a \le x \le b \text{ is an ordered triangle in } X\}$

for any maximal edge $a \le b$ in X. Let $m_{ab} \in |X|$ denote the midpoint of the maximal edge $a \le b$. By assumption, any simplex of X containing a and b has for its vertex set a chain from a to b. Then $B_{|X|}(m_{ab}, \frac{1}{2})$ is a subcomplex of |X| with vertex set $B_{|X|}(m_{ab}, \frac{1}{2}) \cap X^{(0)} = I_{ab}$.

We will prove that Γ is clique-Helly: let $(I_{a_ib_i})_{i \in I}$ denote a family of pairwise intersecting maximal cliques in Γ . Since |X| is injective, there exists $z \in \bigcap_{i \in I} B_{|X|}(m_{a_ib_i}, \frac{1}{2})$. Since each such ball is a subcomplex of |X|, we may assume that z is a vertex of X. We deduce that z belongs to each clique $I_{a_ib_i}$ for $i \in I$. So Γ is clique-Helly.

We will now prove that the triangle complex of Γ is simply connected. Let ℓ denote a combinatorial loop in the 1–skeleton of Γ . Up to homotopy in the triangle complex of Γ , we may assume that ℓ lies in the 1–skeleton of X. Since |X| is injective, it is contractible, so the 2–skeleton of X is simply connected. As the 2-skeleton of X is contained in the triangle complex of Γ , we conclude that ℓ is null-homotopic in the triangle complex of Γ .

According to Theorem 1.12, we deduce that Γ is Helly.

2 The thickening of a lattice

We will explain a very simple construction of Helly graphs and injective metric spaces, starting with a lattice endowed with an action of the group \mathbb{Z} or \mathbb{R} .

Assume that *L* is a lattice such that each upper bounded subset of *L* has a join. Assume that there is an order-preserving, increasing, continuous (with respect to the order topology on *L*) action $(f_t)_{t \in H}$ of $H = \mathbb{Z}$ or \mathbb{R} on *L*, such that,

for all $x, y \in L$, there exists $t \in H_+$ such that $f_{-t}(x) \leq y \leq f_t(x)$.

Let us define the following metric d on L:

for all
$$x, y \in L$$
, $d(x, y) = \inf\{t \in H_+ \mid f_{-t}(x) \le y \le f_t(x)\}$.

Theorem 2.1 • If $H = \mathbb{Z}$, then (L, d) is a Helly graph with the combinatorial distance.

• If $H = \mathbb{R}$, then (L, d) is injective.

Proof We start by proving that *d* is indeed a metric on *L*. If $x, y \in L$ and $t \in H_+$ are such that $f_{-t}(x) \leq y \leq f_t(x)$, then by applying f_t and f_{-t} we deduce that $f_{-t}(y) \leq x \leq f_t(y)$: the metric *d* is symmetric.

We will now prove the triangle inequality: let $x, y, z \in L$, and for $\varepsilon > 0$ consider $t, s \in H_+$ such that $d(x, y) \leq t < d(x, y) + \varepsilon$ and $d(y, z) \leq s < d(y, z) + \varepsilon$. We have

 $f_{-t}(x) \leq y \leq f_t(x)$ and $f_{-s}(y) \leq z \leq f_s(y)$.

Hence

$$f_{-t-s}(x) \leq f_{-s}(y) \leq z \leq f_s(y) \leq f_{s+t}(x),$$

so $d(x, z) \le t + s \le d(x, y) + d(y, z) + 2\varepsilon$. This holds for any $\varepsilon > 0$; hence $d(x, z) \le d(x, y) + d(y, z)$. We will now prove that the metric is positive: assume that $x, y \in L$ are distinct, we will prove that d(x, y) > 0.

• Assume first that x, y are comparable, for instance x < y. Since $\{z \in L \mid z < y\}$ is open and contains x, by continuity of the action there exists $\varepsilon > 0$ such that, for any $|t| \le \varepsilon$, we have $f_t(x) < y$. In particular $d(x, y) \ge \varepsilon$.

• Assume now that x, y are not comparable. Since $U = \{z \in L \mid x \land y < z < x \lor y\}$ is open and contains x, by continuity of the action there exists $\varepsilon > 0$ such that, for any $|t| \le \varepsilon$, we have $f_t(x) \in U$. For any $0 \le t < \varepsilon$, since $x \le f_t(x) < x \lor y$, we have $f_t(x) \not\ge y$. In particular, $d(x, y) \ge \varepsilon$.

Algebraic & Geometric Topology, Volume 24 (2024)

4023

So we have proved that *d* is a metric on *L*. We will now prove that balls in *L* satisfy the Helly property. We will first consider the case $H = \mathbb{Z}$. Let us define a graph \hat{L} with vertex set *L*, and with an edge between $x, y \in L$ if $f_{-1}(x) \leq y \leq f_1(x)$: we will prove that the graph \hat{L} is Helly. We will first prove, by induction on $k \geq 0$, that for any $x \in L$ the ball $B_{\hat{L}}(x, k)$ in the graph \hat{L} coincides with the interval $I(f_{-k}(x), f_k(x)) = \{y \in L \mid f_{-k}(x) \leq y \leq f_k(x)\}.$

For $k \leq 1$ it is the definition of the edges of \hat{L} , so fix $k \geq 2$ and assume that the statement holds for k-1. Fix $y \in I(f_{-k}(x), f_k(x))$, we will prove that $y \in B_{\hat{L}}(x, r)$. Since $y \geq f_{-k}(x)$, we deduce that $f_1(y) \geq f_{-k+1}(x)$, and also since $y \leq f_k(x)$ we deduce that $f_{-1}(y) \leq f_{k-1}(x)$. So we have that both $f_1(y)$ and $f_{k-1}(x)$ are superior to both $f_{-1}(y)$ and $f_{-k+1}(x)$: since L is a lattice, there exists some element $z \in L$ in the intersection $I(f_{-1}(y), f_1(y)) \cap I(f_{-k+1}(x), f_{k-1}(x))$. In particular, y and z are adjacent in \hat{L} , and by induction we know that $d_{\hat{L}}(z, x) \leq k-1$, so $d_{\hat{L}}(x, y) \leq k$. Conversely, it is clear that the ball $B_{\hat{L}}(x, k)$ is included in the interval $I(f_{-k}(x), f_k(x))$. So we have $B_{\hat{L}}(x, k) = I(f_{-k}(x), f_k(x))$. By assumption on the lattice L, we deduce that the graph \hat{L} is connected, and furthermore that $d_{\hat{L}} = d$. Note now that intervals in the lattice L satisfy the Helly property. Fix any collection $(I(x_i, y_i))_{i \in I}$ of

pairwise intersecting intervals of L. Fix $j_0 \in I$. For any $i \in I$, we have $x_i \leq y_{j_0}$, so the set $\{x_i \mid i \in I\}$ is upper bounded. By assumption, the set $\{x_i \mid i \in I\}$ has a join $z \in L$ such that $z \leq y_{j_0}$. This holds for any $j_0 \in I$, so z belongs to the intersection of all intervals $(I(x_i, y_i))_{i \in I}$.

Hence the graph \hat{L} is connected, and its balls satisfy the Helly property: it is a Helly graph.

We now turn to the case $H = \mathbb{R}$, and we will prove that (L, d) is an injective metric space. First note that balls in (L, d) are intervals in L, so according to the previous argument, we know that balls in (L, d) satisfy the Helly property. In order to prove that (L, d) is injective, according to the definition of hyperconvex metric spaces (see for instance [Lang 2013]), it is sufficient to prove that if $x, y \in L$ and $r, s \ge 0$ are such that $d(x, y) \le r + s$, then the balls B(x, r) and B(y, s) intersect. In other words, it is enough to prove that (L, d) is weakly geodesic, ie for any $x, y \in L$ and any $0 \le r \le d(x, y)$, there exists $z \in B(x, r) \cap B(y, d(x, y) - r)$.

For each $k \in \mathbb{N} \setminus \{0\}$, let us consider the action of $\frac{1}{k}\mathbb{Z} \subset \mathbb{R}$ on L, and the associated Helly graph distance,

for all
$$x, y \in L$$
, $d_k(x, y) = \inf \left\{ t \in \frac{1}{k} \mathbb{N} \mid f_{-t}(x) \leq y \leq f_t(x) \right\}.$

Fix $x, y \in L$, and $0 \le r \le d(x, y)$. For each $k \in \mathbb{N} \setminus \{0\}$, there exists $z_k \in L$ such that $d_k(x, z_k) \le r + \frac{1}{k}$ and $d_k(z_k, y) \le d_k(x, y) - r + \frac{1}{k} \le d(x, y) - r + \frac{2}{k}$. So the intervals $I_k = I(f_{-r-1/k}(x), f_{r+1/k}(x))$ and $J_k = I(f_{-d(x,y)+r-2/k}(y), f_{d(x,y)-r+2/k}(y))$ in L intersect. If $k \le k'$, we know that $I_{k'} \subset I_k$ and $J_{k'} \subset J_k$. We deduce that the family of intervals $\{I_k\}_{k\in\mathbb{N}\setminus\{0\}} \cup \{J_k\}_{k\in\mathbb{N}\setminus\{0\}}$ pairwise intersects. By the Helly property for intervals, the global intersection is nonempty: let us denote by z some element in the intersection. We know that $\lim_{k\to+\infty} d_k(x,z) = d(x,z) \le r$ and $\lim_{k\to+\infty} d_k(y,z) = d(y,z) \le d(x,y) - r$, so we have proved that (L,d) is a weakly geodesic metric space. Since balls in (L,d) satisfy the Helly property, we conclude that (L,d) is injective.

An immediate consequence concerns Garside groups (see [Dehornoy et al. 2015; McCammond 2006; Haettel and Huang 2024]). Recall that a group *G* is called *Garside* if there exists a subset $S \subset G$ and an element $\delta \in G$ such that the following hold:

- S spans the group G.
- For each element in the semigroup $\langle S \rangle^+$, there is a bound on the length of factorizations over S.
- The element δ belongs to the semigroup (S)⁺, and δ is *balanced*: the set of prefixes of δ coincides with the set of suffixes of δ.
- The poset of prefixes of δ in $\langle S \rangle^+$ is a lattice.

Note that G is endowed with two natural orders, the left order \leq_L and the right order \leq_R :

- $g \leq_L h$ if and only if $g^{-1}h \in \langle S \rangle^+$.
- $g \leq_R h$ if and only if $hg^{-1} \in \langle S \rangle^+$.

Authors sometimes add the requirement that *S* is finite, in which case *G* may also be called a Garside group of finite type. Also note that, given a Garside group (G, S, δ) , for any $g \in G$, there exists $n \in \mathbb{N}$ such that $\delta^{-n} \leq_L g \leq_L \delta^n$.

Fix a Garside group (G, S, δ) . Let X denote the graph with vertex set G, with an edge between $g, h \in G$ if $g\delta^{-1} \leq_L h \leq_L g\delta$. The graph X is called the *thickening* of G. In relation to Definition 1.7, it corresponds to the cell complex with vertex set G, whose maximal cells are translates $(g[\Delta^{-1}, \Delta]_{\leq_L})_{g \in G}$ of the interval $[\Delta^{-1}, \Delta]$.

Corollary 2.2 The thickening of any Garside group is a Helly graph.

Proof The left order on *G* is a lattice order (see [Dehornoy et al. 2015]). Furthermore, consider the action of \mathbb{Z} on *G* by right multiplication by δ . For any $g \in G$, there exists $n \in \mathbb{N}$ such that $\delta^{-n} \leq_L g \leq_L \delta^n$. So this action satisfies the assumptions of Theorem 2.1: we deduce that the graph *X* is Helly. \Box

Note that this applies, in particular, to Garside groups of infinite type, such as crystallographic Garside groups (see [McCammond and Sulway 2017]). However, we do not have yet an application of this simply transitive action of a Garside group on a locally infinite Helly graph.

In the case of Garside groups of finite type, we recover a particularly simple proof of the following result by Huang and Osajda [2021]. In particular, our proof does not rely on the deep local-to-global result for Helly graphs (Theorem 1.12, see [Chalopin et al. 2020b]).

Corollary 2.3 (Huang–Osajda) Any Garside group of finite type is a Helly group.

In particular, this leads to a particularly simple proof that braid groups are Helly, relying only on some Garside structure.

3 The affine version of a lattice

In this section, we will prove Theorem 3.10 stating that the orthoscheme complex of a bounded graded lattice, endowed with the orthoscheme ℓ^{∞} metric, is injective. In order to do so, we will apply results from the previous section, and endow the geometric realization of a lattice with a partial order, which is a lattice.

Assume that *L* is a bounded, graded lattice of rank *n*. Let \leq_L denote the order on *L*. Let *H* denote either a cyclic subgroup of $(\mathbb{R}, +)$ or $H = \mathbb{R}$. We will define a new poset M_H , which will be called the *affine version* of *L* over *H*. If there is no ambiguity about *H*, we will simply write $M = M_H$. Let C(L) denote the set of maximal chains $c_{0,1} = 0 <_L c_{1,2} <_L \cdots <_L c_{n-1,n} <_L c_{n,n+1} = 1$ in *L*. We will use the convention that the element denoted by $c_{i,i+1}$ has rank *i*.

Let us consider the subspace

$$\sigma = \{ u \in H^n \mid u_1 \leq u_2 \leq \ldots \leq u_n \}$$

of H^n .

For each maximal chain $c \in C(L)$, let σ_c denote a copy of σ .

Let us consider the space

$$M = \bigcup_{c \in C(L)} \sigma_c / \sim,$$

where for each $c, c' \in C(L)$, if we let $I = \{1 \le i \le n-1 \mid c_{i,i+1} \ne c'_{i,i+1}\}$, we identify σ_c and $\sigma_{c'}$ along the subspaces

$$\{u \in \sigma_c \mid \text{for all } i \in I, u_i = u_{i+1}\} \simeq \{u \in \sigma_{c'} \mid \text{for all } i \in I, u_i = u_{i+1}\}$$

We can describe the set of elements of M as a quotient of the space $M_0 = C(L) \times \sigma$.

Example One illustrating example is the following: consider the Boolean lattice L of rank n, ie the lattice of subsets of the finite set $\{1, \ldots, n\}$, with the inclusion order. Maximal chains in L correspond to permutations of $\{1, \ldots, n\}$. The space M_H may be identified with H^n , where, for each permutation w of $\{1, \ldots, n\}$, the subspace σ_w is

$$\sigma_w = \{ x \in H^n \mid x_{w(1)} \leq x_{w(2)} \leq \ldots \leq x_{w(n)} \}.$$

If $c \in C(L)$ and $u \in \sigma$, let us denote by [c, u] the equivalence class of $(c, u) \in M_0$ in M.

For each $c \in C(L)$, let us endow σ_c with the partial order from $H^n \subset \mathbb{R}^n$: $u \leq v$ if, for all $1 \leq i \leq n$, $u_i \leq v_i$. Let us endow M with the induced partial order: we have $\alpha \leq \beta$ in M if there exists a sequence $\alpha_0 = \alpha, \alpha_1, \ldots, \alpha_m = \beta$ in M such that, for each $0 \leq i \leq m - 1$, there exists $c \in C(L)$ and $x \leq y$ in σ_c such that $\alpha_i = [c, x]$ and $\alpha_{i+1} = [c, y]$.

Let \leq_M denote the order on *M*. We will prove the following.

Theorem 3.1 If H is a discrete subgroup of \mathbb{R} , the poset M_H is a lattice.

Before proving Theorem 3.1, we will gather some preliminary results. Without loss of generality, assume that $H = \mathbb{Z}$. To simplify notation, we will let $M = M_H$.

First notice that if $(a, u) \sim (b, v)$, then u = v. Therefore the second projection $M_0 \rightarrow \sigma$ defines a projection $\pi: M \rightarrow \sigma$. Fix $\beta = [b, v], \gamma = [c, w] \in M$, and fix $\alpha \leq_M \beta, \gamma$ in M. We will prove that β, γ have a join, and α will play an auxiliary role.

We say that β is *elementarily superior* to α if there are representatives $\alpha = [a, u]$ and $\beta = [b, v]$ such that a = b and there exist $1 \le i \le j \le n$ such that

- $u_i = u_{i+1} = \cdots = u_j$,
- $v_i = v_{i+1} = \dots = v_j = u_j + 1$, and
- for all $k \notin [i, j]$, $u_k = v_k$.

Lemma 3.2 Fix $\alpha = [a, u] \in M$. Any element $\beta = [b, v]$ of M elementarily superior to α is uniquely determined by:

- integers $1 \le i \le j \le n$, such that $u_i = u_{i+1} = \cdots = u_j$, and $u_j < u_{j+1}$ if j < n,
- some element $b_{i-1,i}$ of rank i-1 in the interval $I(a_{i_0-1,i_0}, a_{j,j+1})$, where $i_0 \in [1, i]$ is minimal such that $u_{i_0} = u_i$.

Let us set $\beta = \alpha[i, j, b_{i-1,i}]$.

Proof Let us define the element $\beta = [b, v]$ in M by

for all $k \notin [i, j]$,	$v_k = u_k,$	for all $k \leq i_0 - 1$,	$b_{k,k+1} = a_{k,k+1},$
for all $k \in [i, j]$,	$v_k = u_j + 1,$	for all $k \ge j$,	$b_{k,k+1} = a_{k,k+1}.$

Note that v is nondecreasing; hence $v \in \sigma$. Furthermore, since $v_{i_0} = v_{i_0+1} = \cdots = v_{i-1}$ and $v_i = v_{i+1} = \cdots = u_j + 1 \leq v_{j+1}$, it is enough to define $b_{k,k+1}$ for $k \leq i_0 - 1$, k = i - 1 and $k \geq j$. Hence β is well-defined, and it is elementarily superior to α . It is clear that β is the only such element in M. \Box

Lemma 3.3 Given $\alpha \leq_M \beta$ in M, there exist $m \geq 0$ and a sequence $\beta_0 = \alpha, \beta_1, \dots, \beta_m = \beta$ for which, for each $0 \leq i \leq m - 1$, β_{i+1} is elementarily superior to β_i .

Proof According to the definition of the order on M, it is sufficient to prove the statement for $\alpha = [a, u]$ and $\beta = [b, v]$ such that a = b. We have $u \leq v$ in $\mathbb{Z}^n \subset \mathbb{R}^n$. Then consider a sequence $u^0 = u \leq u^1 \leq \ldots \leq u^m = v$ such that, for each $0 \leq k \leq m-1$, there exist $1 \leq i_k \leq j_k \leq n$ such that

• $u_{i_k}^k = u_{i_k+1}^k = \dots = u_{j_k}^k$,

•
$$u_{i_k}^{k+1} = u_{i_k+1}^{k+1} = \dots = u_{j_k}^{k+1} = u_{j_k}^k + 1$$
, and

• for all $\ell \notin [i_k, j_k], \ u_\ell^k = u_\ell^{k+1}$.

For each $0 \le k \le m-1$, the element $\beta_{k+1} = [a, u^{k+1}]$ is elementarily superior to $\beta_k = [a, u^k]$. \Box

Algebraic & Geometric Topology, Volume 24 (2024)

Lemma 3.4 We have $\alpha = [a, u] \leq \beta = [b, v]$ in M if and only if $u \leq v$ and for every $0 \leq j \leq n-1$ such that $u_j < u_{j+1}$. If we denote by $i \in \{0, ..., j\}$ the minimal element such that $v_{i+1} \geq u_{j+1}$, we have $b_{i,i+1} \leq L a_{j,j+1}$.

Proof Let us denote by \prec this relation. We first show that \prec is transitive and antisymmetric.

Assume that $\alpha = [a, u] \leq \beta = [b, v]$ and u = v; we will prove that $\alpha = \beta$. Fix $0 \leq j \leq n-1$, and assume that $a_{j,j+1} \neq b_{j,j+1}$. Since these two elements of *L* have the same rank *j*, we deduce that they are not comparable: $b_{j,j+1} \not\leq L a_{j,j+1}$ and $a_{j,j+1} \not\leq L b_{j,j+1}$. We will prove that $u_j = u_{j+1}$: by contradiction, if $u_j < u_{j+1}$, then the minimal $i \in \{0, ..., j\}$ such that $v_{i+1} \geq u_{j+1}$ is equal to *j*, so $b_{j,j+1} \leq_L a_{j,j+1}$. So $u_j = u_{j+1}$. This proves that $\alpha = \beta$.

This implies in particular that \prec is antisymmetric: indeed assume that $\alpha \preccurlyeq \beta$ and $\beta \preccurlyeq \alpha$. Then $u \leqslant v$ and $v \leqslant u$, so u = v; hence $\alpha = \beta$.

We now prove that \prec is transitive: assume that $\alpha = [a, u] \preccurlyeq \beta = [b, v] \preccurlyeq \gamma = [c, w]$. Then $u \leqslant v \leqslant w$, so $u \leqslant w$. Let $0 \leqslant k \leqslant n-1$ such that $u_k < u_{k+1}$, and let $i \in \{0, \dots, k\}$ be minimal such that $w_{i+1} \ge u_{k+1}$: we want to prove that $c_{i,i+1} \leqslant_L a_{k,k+1}$. Let $j \in \{0, \dots, k\}$ be minimal such that $v_{j+1} \ge u_{k+1}$: we know that $b_{j,j+1} \leqslant_L a_{k,k+1}$. Since j is minimal, we know that either j = 0 or $v_j < u_{k+1}$.

If j = 0, then since $i \leq j$ we know that i = 0 and so $c_{i,i+1} = 0 = b_{j,j+1} \leq a_{k,k+1}$.

If $v_j < u_{k+1}$ then $v_j < v_{j+1}$, and let $i' \in \{0, \dots, j\}$ be minimal such that $w_{i'+1} \ge v_{j+1}$. We know that $c_{i',i'+1} \le L b_{j,j+1}$. But since $w_i < u_{k+1} \le v_{j+1}$, we have $i \le i'$, so $c_{i,i+1} \le L c_{i',i'+1} \le L b_{j,j+1} \le L a_{k,k+1}$. Hence $\alpha \le \gamma$.

We will now prove that if $\beta = [b, v]$ is elementarily superior to $\alpha = [a, u]$, we have $\alpha \preccurlyeq \beta$. Let us set $\beta = \alpha[i, j, b_{i-1,i}]$. First note that $u \leqslant v$. If $0 \leqslant k \leqslant n-1$ is such that $u_k < u_{k+1}$, then either $k \leqslant i-1$ or $k \ge j$. Let $\ell \in \{0, \dots, k\}$ denote the minimal element such that $v_{\ell+1} \ge u_{k+1}$; we will prove that $b_{\ell,\ell+1} \leqslant L a_{k,k+1}$.

- If $k \leq i-1$, then $\ell = k$ and $b_{\ell,\ell+1} = a_{k,k+1}$.
- If $k \ge j$ and $v_{k+1} = v_j$, then $\ell = i 1$ and $b_{\ell,\ell+1} = b_{i-1,i} \le_L a_{j,j+1} \le_L a_{k,k+1}$.
- If $k \ge j$ and $v_{k+1} > v_j$, then $\ell \ge j$ and $b_{\ell,\ell+1} = a_{\ell,\ell+1} \le_L a_{k,k+1}$.

According to Lemma 3.3, we deduce that if $\alpha \leq_M \beta$, then $\alpha \leq \beta$.

We will now prove that if $\alpha = [a, u] \leq \beta = [b, v]$ with $\alpha \neq \beta$, there exists γ elementarily superior to α such that $\gamma \leq \beta$. If u = v, we have seen that $\alpha = \beta$, so let us assume that u < v. Let $1 \leq i \leq n$ be minimal such that $u_i < v_i$. Let $j \in \{i, ..., n\}$ be maximal such that $u_i = u_j$.

Let $i_0 \in \{1, ..., i\}$ be minimal such that $u_{i_0} = u_i$. Since for all $k \leq i - 1$ we have $u_k = v_k$, there exist representatives $\alpha = [a, u]$ and $\beta = [b, v]$ such that, for any $1 \leq k \leq i - 1$, we have $a_{k-1,k} = b_{k-1,k}$.

We will also prove that we may assume that $a_{i-1,i} = b_{i-1,i}$.

Assume first that $u_{i-1} < u_i$. Let $p \in \{0, ..., i-1\}$ be minimal such that $v_{p+1} \ge u_i$. Since $v_{i-1} = u_{i-1} < u_i$, we deduce that p = i - 1. Since $\alpha \le \beta$, we have $b_{i-1,i} \le L a_{i-1,i}$. As both elements have the same rank i - 1 in L, we have $a_{i-1,i} = b_{i-1,i}$.

Assume now that $u_{i-1} = u_i$. In the case j < n, we have $u_j < u_{j+1}$: let $p \in \{0, \dots, j\}$ be minimal such that $v_{p+1} \ge u_{j+1}$. Since $v_{i-1} = u_{i-1} < u_{j+1}$, we deduce that $p \ge i-1$. Since $\alpha \le \beta$, we have $b_{i-1,i} \le L b_{p,p+1} \le L a_{j,j+1}$. Since $u_{i-1} = u_i = \dots = u_j$, we may choose a representative of α such that $a_{i-1,i} = b_{i-1,i}$. In the case j = n, we have $u_{i-1} = u_i = \dots = u_n$; we may also choose a representative of α such that $a_{i-1,i} = b_{i-1,i}$.

We have proved that we may assume that $a_{i-1,i} = b_{i-1,i}$.

In the case j = n, since $u_i = \cdots = u_n$, we may choose a representative of α such that a = b. In this case, the element $\gamma = \alpha[i, n, b_{i-1,i}]$ is elementarily superior to α , and $\gamma \leq \beta$.

For the rest of the proof, assume that j < n.

We know that $b_{i-1,i} \ge_L b_{i_0-1,i_0} = a_{i_0-1,i_0}$. Since $u_j < u_{j+1}$ and $\alpha \le \beta$, if we denote by $i' \in \{0, \ldots, j\}$ the minimal element such that $v_{i'+1} \ge u_{j+1}$, we have $b_{i',i'+1} \le_L a_{j,j+1}$. Since $v_{i-1} = u_{i-1} < u_{j+1}$, we have i' > i-2, so $b_{i-1,i} \le_L b_{i',i'+1} \le_L a_{j,j+1}$.

So we know that $b_{i-1,i} \in I(a_{i_0-1,i_0}, a_{j,j+1})$: we can define $\gamma = \alpha[i, j, b_{i-1,i}] = [w, c]$. We will now prove that $\gamma \preccurlyeq \beta$. Fix $0 \le k \le n-1$ such that $w_k < w_{k+1}$, and denote by $\ell \in \{0, \dots, k\}$ the minimal element such that $v_{\ell+1} \ge w_{k+1}$; we will prove that $b_{\ell,\ell+1} \le L c_{k,k+1}$. Recall that either $k \le i-1$ or $k \ge j$.

- If $k \leq i-2$, then $w_{k+1} = u_{k+1}$. Since $\alpha \leq \beta$, we know that $b_{\ell,\ell+1} \leq u_{k,k+1} = c_{k,k+1}$.
- If k = i 1, then $w_{k+1} = w_i = u_i + 1 > u_i$. Then $v_{i-1} = u_{i-1} \le u_i < w_{k+1}$, so $\ell = i 1 = k$. Hence $b_{\ell,\ell+1} = b_{i-1,i} = a_{i-1,i} = c_{i-1,i} = c_{k,k+1}$.
- If $k \ge j$, then $w_{k+1} = u_{k+1}$. Since $\alpha \le \beta$, we know that $b_{\ell,\ell+1} \le a_{k,k+1}$. We also have $a_{k,k+1} = c_{k,k+1}$, so $b_{\ell,\ell+1} \le c_{k,k+1}$.

So we conclude that $\gamma \preccurlyeq \beta$.

Now fix any $\alpha = [a, u] \preccurlyeq \beta = [b, v]$. By induction on $||v - u||_1$, we see that there is a bound on sequences of elementarily superior elements starting from α which all are $\preccurlyeq \beta$. Therefore we conclude that $\alpha \leq_M \beta$. In conclusion, the two orders \leq_M and \prec coincide.

Given $\alpha \leq_M \beta$ in *M*, let us denote by $D(\alpha, \beta)$ the minimal number $m \geq 0$ such that there exists a sequence $\beta_0 = \alpha, \beta_1, \dots, \beta_m = \beta$ for which, for each $0 \leq i \leq m-1$, β_{i+1} is elementarily superior to β_i . We will prove, by induction on $D(\alpha, \beta) + D(\alpha, \gamma)$, that β and γ have a join.

Lemma 3.5 Assume that $\alpha, \beta, \gamma \in M$ are such that $\alpha \leq_M \beta$, $\alpha \leq_M \gamma$ and $D(\alpha, \beta) = D(\alpha, \gamma) = 1$. Then β, γ have a join δ such that $D(\beta, \delta) \leq 1$ and $D(\gamma, \delta) \leq 1$.

Proof Consider representatives $\alpha = [a, u]$, $\beta = [b, v]$ and $\gamma = [c, w]$ of α , β and γ respectively. According to Lemma 3.2, there exist $1 \le i \le j \le n$ and $b_{i,i+1}$ such that $\beta = \alpha[i, j, b_{i,i+1}]$, and there exist $1 \le i' \le j' \le n$ and $c_{i',i'+1}$ such that $\gamma = \alpha[i', j', c_{i',i'+1}]$.

First case Assume that the intervals [i, j] and [i', j'] are disjoint, for instance j < i'. Let us define $\delta = \beta[i', j', c_{i',i'+1}]$: we will see that δ is well-defined and that $\delta = \gamma[i, j, b_{i,i+1}]$.

First note that $v_{j'} = u_{j'} < u_{j'+1} = v_{j'+1}$. Furthermore, let $i'_0 \in [1, i']$ be minimal such that $u_{i'_0} = u_{i'}$. Since $u_j < u_{j+1}$, we know that $j + 1 \le i'_0$. So we have $v_{i'_0} = u_{i'_0} = u_{i'} = v_{i'}$. So, if we denote by $i''_0 \in [1, i']$ the minimal integer such that $v_{i''_0} = v_{i'}$, we have $i''_0 \le i'_0$.

By definition we have $c_{i',i'+1} \in I(a_{i'_0-1,i'_0}, a_{j',j'+1})$. Since j < j', we have $b_{j',j'+1} = a_{j',j'+1}$, so $c_{i',i'+1} \leq_L b_{j',j'+1}$. And as $b_{i''_0-1,i''_0} \leq_L b_{i'_0-1,i'_0} = a_{i'_0-1,i'_0}$, we deduce that $c_{i',i'+1} \geq_L a_{i'_0-1,i'_0} \geq_L b_{i''_0-1,i''_0}$. So we have $c_{i',i'+1} \in I(b_{i''_0-1,i''_0}, b_{j',j'+1})$. Hence $\delta = \beta[i', j', c_{i',i'+1}]$ is well-defined, and it is elementarily superior to β .

Following the same argument, the element $\delta' = \gamma[i, j, b_{i,i+1}]$ is well-defined. We will prove that $\delta = \delta'$. Let $i_0 \in [1, i]$ be minimal such that $u_{i_0} = u_i$. According to the proof of Lemma 3.2, we can see that $\delta = \delta' = [d, x]$ are explicitly equal to the following:

for all
$$k \notin [i, j] \cup [i', j']$$
, $x_k = u_k$,
for all $k \in [i, j]$, $x_k = u_j + 1$,
for all $k \in [i', j']$, $x_k = u_{j'} + 1$,
for all $k \notin i_0 - 1$, $d_{k,k+1} = a_{k,k+1}$,
 $d_{i,i+1} = b_{i,i+1}$,
for all $k \notin [j, i'_0 - 1]$, $d_{k,k+1} = a_{k,k+1}$,
 $d_{i',i'+1} = c_{i',i'+1}$,
for all $k \geqslant j'$, $d_{k,k+1} = a_{k,k+1}$.

So the element $\delta = \delta'$ is elementarily superior to both β and γ .

We will now prove that δ is the minimal element of M superior to both β and γ . Fix $\theta = [e, y] \in M$ as any element superior to both β and γ ; we will prove that $\delta \leq \theta$. Since $y \geq v$ and $y \geq w$, we deduce that $y \geq x$. Fix any $0 \leq k \leq n-1$ such that $x_k < x_{k+1}$, and let $\ell \in \{0, \dots, k\}$ denote the minimal element such that $y_{\ell+1} \geq x_{k+1}$. We will prove that $e_{\ell,\ell+1} \leq L d_{k,k+1}$.

- Assume that $k \leq j-1$. Then $x_{k+1} = v_{k+1}$, and since $\beta \leq \theta$, we deduce by Lemma 3.4 that $e_{\ell,\ell+1} \leq_L b_{k,k+1} = d_{k,k+1}$.
- Assume that $k \ge j$. Then $x_{k+1} = w_{k+1}$, and since $\gamma \le \theta$, we deduce by Lemma 3.4 that $e_{\ell,\ell+1} \le c_{k,k+1} = d_{k,k+1}$.

According to Lemma 3.4, we deduce that $\delta \leq_M \theta$: δ is the minimal element of M superior to both β and γ . Hence $\delta = \beta \vee_M \gamma$. Furthermore, we have noticed that δ is elementarily superior to both β and γ , so $D(\beta, \delta) = D(\gamma, \delta) = 1$.

Second case Assume now that the intervals [i, j] and [i', j'] intersect. Without loss of generality, assume that $i \leq i'$. Since $u_j < u_{j+1}$, we deduce that j = j'. Let $1 \leq i_0 \leq n$ be minimal such that $u_{i_0} = u_i$. The elements $b_{i-1,i}$ and $c_{i'-1,i'}$ both belong to the interval $I(a_{i_0-1,i_0}, a_{j,j+1})$.

If $b_{i-1,i} = c_{i'-1,i'}$, then $\beta = \gamma$ and they have a trivial join $\delta = \beta = \gamma$. So we may assume that $b_{i-1,i} \neq c_{i'-1,i'}$.

If $b_{i-1,i} <_L c_{i'-1,i'}$, we have $\beta \leq \gamma$, so β and γ have a join $\delta = \gamma$ which satisfies $D(\beta, \delta) = 1$ and $D(\gamma, \delta) = 0$. Let us assume now that $b_{i-1,i} \not\leq_L c_{i'-1,i'}$.

Consider the meet $g = b_{i-1,i} \wedge_L c_{i'-1,i'} \in L$: its rank r-1 is such that $i_0 \leq r < i, i'$. Let us define $\delta = \alpha[r, j, g] \in M$. We see that $\delta = \beta[r, i-1, g] = \gamma[r, i'-1, g]$, so δ is elementarily superior to β and γ .

We will now prove that $\delta = [d, x]$ is the minimal element of M superior to both β and γ . Fix $\theta = [e, y] \in M$ as any element superior to both β and γ ; we will prove that $\delta \leq \theta$.

We will first prove that $x \leq y$. Since $\beta, \gamma \leq_M \theta$, we deduce that $v, w \leq y$. In particular, for any m < r or $m \geq i$, we have $y_m \geq b_m = x_m$. And for $r \leq m \leq i-1$, we have $y_m \geq a_m = x_m - 1$. Assume by contradiction that there exists $m \in \{r, \ldots, i-1\}$ such that $y_m = x_m - 1$, and choose such m maximal. Since $x_r = x_{r+1} = \cdots = x_j$, we have $y_{m+1} \geq x_{m+1} = x_i = v_i = x_{i'} = w_{i'}$. Since $\beta \leq_L \theta$ and $\gamma \leq_L \theta$, according to Lemma 3.4, we know that $e_{m,m+1} \leq_L b_{i-1,i}$ and $e_{m,m+1} \leq_L c_{i'-1,i'}$. In particular, we deduce that $e_{m,m+1} \leq_L b_{i-1,i} \wedge_L c_{i'-1,i'} = g$. Note that the rank of $e_{m,m+1}$ is m, whereas the rank of g is r-1. Since m > r-1, this is a contradiction. Hence $x \leq y$.

Fix any $0 \le k \le n-1$ such that $x_k < x_{k+1}$, and let $\ell \in \{0, \dots, k\}$ denote the minimal element such that $y_{\ell+1} \ge x_{k+1}$. We will prove that $e_{\ell,\ell+1} \le L d_{k,k+1}$.

- Assume that $k \leq r-2$. Then $x_{k+1} = v_{k+1}$, and since $\beta \leq \theta$ we deduce by Lemma 3.4 that $e_{\ell,\ell+1} \leq_L b_{k,k+1} = d_{k,k+1}$.
- Assume that $r-1 \le k \le j-1$. Then $x_{k+1} = v_i = w_{i'}$. Since $\beta \le \theta$, we deduce by Lemma 3.4 that $e_{\ell,\ell+1} \le L b_{i-1,i}$. And since $\gamma \le \theta$, we also deduce that $e_{\ell,\ell+1} \le L c_{i'-1,i'}$. Hence $e_{\ell,\ell+1} \le L b_{i-1,i} \land L c_{i'-1,i'} = g = d_{r-1,r} \le L d_{k,k+1}$.
- Assume that $k \ge j$. Then $x_{k+1} = v_{k+1}$, and since $\beta \le \theta$ we deduce by Lemma 3.4 that $e_{\ell,\ell+1} \le b_{k,k+1} = d_{k,k+1}$.

According to Lemma 3.4, we deduce that $\delta \leq_M \theta$: δ is the minimal element of M superior to both β and γ . Hence $\delta = \beta \lor_M \gamma$. Furthermore, we have noticed that δ is elementarily superior to both β and γ , so $D(\beta, \delta) = D(\gamma, \delta) = 1$.

Lemma 3.6 Assume that $\alpha, \beta, \gamma \in M$ are such that $\alpha \leq_M \beta$, $\alpha \leq_M \gamma$, $D(\alpha, \beta) = m$ and $D(\alpha, \gamma) = m'$ for some $m, m' \in \mathbb{N}$. Then β, γ have a meet δ such that $D(\beta, \delta) \leq m'$ and $D(\gamma, \delta) \leq m$.

Proof We proceed by induction on m + m': when $m + m' \leq 2$, the statement holds by Lemma 3.5. Now fix $k \geq 3$, and assume that the statement holds when m + m' < k. Fix m, m' such that m + m' = k, and without loss of generality assume that $m \geq 2$. Choose $\beta_0 = \alpha, \beta_1, \ldots, \beta_m = \beta$ an elementary sequence from α to β , with $m = D(\alpha, \beta)$. We have $D(\alpha, \beta_1) + D(\alpha, \gamma) = 1 + m' < k$, so by induction there exists $\delta' = \beta_1 \vee_M \gamma$ with $D(\beta_1, \delta') \leq m'$ and $D(\gamma, \delta') \leq 1$. Since $D(\beta_1, \beta) + D(\beta_1, \delta') \leq m - 1 + m' < k$, by induction there exists $\delta = \beta \vee_M \delta'$ such that $D(\beta, \delta) \leq m'$ and $D(\delta', \delta) \leq m - 1$. So we deduce that $D(\gamma, \delta) \leq m$.

We will now prove that δ is the meet of β and γ . We have $\delta \ge_M \beta$ and $\delta \ge_M \delta' \ge_M \gamma$. Furthermore, consider any $\theta \in M$ such that $\theta \ge_M \beta$ and $\theta \ge_M \gamma$. As $\beta \ge_M \beta_1$, we deduce that $\theta \ge_M \beta_1 \lor_M \gamma = \delta'$. And we deduce that $\theta \ge_M \beta \lor_M \delta' = \delta$. So we have proved that $\delta = \beta \lor_M \gamma$. \Box

Proof of Theorem 3.1 Fix any $\beta = [b, v]$, $\gamma = [c, w] \in M$. Let $k \ge 0$ such that $v_n - k < w_1$. Let $u = (v_1 - k, v_2 - k, \dots, v_n - k)$. Then $\alpha = [b, u] \in M$ is inferior to β , and we will see that it is also inferior to γ . Indeed let $\gamma' = [c, w']$, where $w' = (w_1, w_1, \dots, w_1)$. Since $w' \le w$, we have $\gamma' \le_M \gamma$. On the other hand, since $\gamma' = [b, w']$ and $u \le w'$, we have $\alpha \le_M \gamma'$, so $\alpha \le_M \gamma$.

We can now apply Lemma 3.6 to deduce that β and γ have a meet in M. By symmetry of the construction, β and γ also have a join in M. So M is a lattice.

If $H = \mathbb{R}$, the affine version $M_{\mathbb{R}}$ of L over \mathbb{R} is a gluing of subspaces $\sigma \subset \mathbb{R}^n$. We may therefore endow $M_{\mathbb{R}}$ with the piecewise length metric $d_{\mathbb{R}}$ induced by the standard ℓ^{∞} metric on each $\sigma \subset \mathbb{R}^n$.

Let us define an action of \mathbb{R} on $M_{\mathbb{R}}$ as

$$\mathbb{R} \times M_{\mathbb{R}} \to M_{\mathbb{R}}, \quad (t, [a, u]) \mapsto t \cdot [a, u] = [a, (u_1 + t, u_2 + t, \dots, u_n + t)].$$

This action is well-defined, preserves the order \leq_M , is increasing and continuous. Moreover, we have the following property.

Lemma 3.7 For any $\alpha, \beta \in M_{\mathbb{R}}$, there exists t > 0 such that $(-t) \cdot \alpha \leq_M \beta \leq_M t \cdot \alpha$.

Proof Consider representatives $\alpha = [a, u]$ and $\beta = [b, v]$ of α and β respectively. Let t > 0 such that $v_n \leq u_1 + t$ and $u_n \leq v_1 + t$. Then, if we let $\gamma = [a, (v_n, \dots, v_n)] = [b, (v_n, \dots, v_n)]$, we have $\beta \leq_M \gamma \leq_M t \cdot \alpha$; hence $\beta \leq_M t \cdot \alpha$. Similarly, we have $(-t) \cdot \alpha \leq_M \beta$.

Theorem 3.8 If $H = \mathbb{R}$, the poset $M_{\mathbb{R}}$ is a lattice, and the metric space $(M_{\mathbb{R}}, d_{\mathbb{R}})$ is injective.

Proof Note that, for each $\theta > 0$, the space $M_{\theta\mathbb{Z}}$ may be realized naturally as a closed subspace of $M_{\mathbb{R}}$. Furthermore, the sequence of closed subsets $M_{\theta\mathbb{Z}}$ of $M_{\mathbb{R}}$ converges to $M_{\mathbb{R}}$ as $\theta \to 0$. We will use this

convergence to prove that $M_{\mathbb{R}}$ is a lattice. We will then prove that the assumptions of Theorem 2.1 are satisfied.

We will now prove that any pair of elements in $M_{\mathbb{R}}$ have a join. Fix $\alpha, \beta \in M_{\mathbb{R}}$; we will define a common upper bound γ for α and β .

Consider maximal chains a, b in L such that $\alpha \in \sigma_a$ and $\beta \in \sigma_b$. For any $\theta > 0$, note that, for any $x \in \mathbb{R}$, there exists $x_{\theta} \in \theta\mathbb{Z}$ such that $x_{\theta} - \theta \leq x \leq x_{\theta} + \theta$. So we may consider $\alpha_{\theta} \in M_{\theta\mathbb{Z}} \cap \sigma_a$ and $\beta_{\theta} \in M_{\theta\mathbb{Z}} \cap \sigma_b$ such that $(-\theta) \cdot \alpha_{\theta} \leq_M \alpha \leq_M \theta \cdot \alpha_{\theta}$, and similarly $(-\theta) \cdot \beta_{\theta} \leq_M \beta \leq_M \theta \cdot \beta_{\theta}$.

According to Theorem 3.1, the poset $M_{\theta\mathbb{Z}}$ is a lattice: consider $\gamma_{\theta} = \alpha_{\theta} \vee_{M_{\theta\mathbb{Z}}} \beta_{\theta}$. Let $C_{a,b} \subset L$ denote the smallest subset of L containing a, b, and which is stable under meets. Since L is bounded and graded, $C_{a,b}$ is finite. According to the proof of Theorem 3.1, we see that, for every $\theta > 0$, there exists a maximal chain $c_{\theta} \subset C_{a,b}$ such that $\gamma_{\theta} \in \sigma_{c_{\theta}}$. Since $C_{a,b}$ is finite, σ is locally compact and $(\gamma_{\theta})_{\theta>0}$ is bounded, there exists a sequence $\theta_k \xrightarrow{k \to +\infty} 0$ such that the sequence $(\gamma_{\theta_k})_{k \in \mathbb{N}}$ converges to some $\gamma \in M_{\mathbb{R}}$. Note that, for any $\theta > 0$, we have $\gamma_{\theta} \ge_M \alpha_{\theta} \ge_M (-\theta) \cdot \alpha$. Since the sequence $(\gamma_{\theta_k})_{k \in \mathbb{N}}$ converges to γ , and the sequence $((-\theta_k) \cdot \alpha)_{k \in \mathbb{N}}$ converges to α by continuity of the action, we deduce that $\gamma \ge_M \alpha$. Similarly $\gamma \ge_M \beta$.

So γ is a common upper bound for α and β . We will now prove that γ is a minimal upper bound, which will prove that γ is the join of α and β . Let us consider an upper bound $\delta \in M_{\mathbb{R}}$ of α and β ; we will prove that $\gamma \leq_M \delta$. For any $\theta > 0$, fix $\delta_{\theta} \in M_{\theta\mathbb{Z}}$ such that $(-\theta) \cdot \delta_{\theta} \leq_M \delta \leq_M \theta \cdot \delta_{\theta}$. In particular $(2\theta) \cdot \delta_{\theta} \geq_M \theta \cdot \delta \geq_M \theta \cdot \alpha \geq_M \alpha_{\theta}$, and similarly $(2\theta) \cdot \delta_{\theta} \geq_M \beta_{\theta}$. We deduce that $(2\theta) \cdot \delta_{\theta} \geq_M \alpha_{\theta} \vee_{M_{\theta\mathbb{Z}}} \beta_{\theta} = \gamma_{\theta}$. Considering the limit along $(\theta_k)_{k \in \mathbb{N}}$ as $k \to +\infty$, we deduce that $\delta \geq_M \gamma$. So we have proved that α and β have a join γ in $M_{\mathbb{R}}$. By symmetry of the construction, they also have a meet, so $M_{\mathbb{R}}$ is a lattice.

We now turn to the assumptions of Theorem 2.1: we will first prove that every upper bounded subset of $M_{\mathbb{R}}$ has a join. Since $M_{\mathbb{R}}$ is a lattice, it is enough to prove that every bounded, increasing sequence is convergent. Fix an increasing sequence $(\alpha_k)_{k\geq 0}$ in $M_{\mathbb{R}}$, bounded above by some $\alpha \in M_{\mathbb{R}}$.

We will prove an intermediate result concerning ℓ^1 metrics. Let us endow $M_{\mathbb{R}}$ with the length metric d_1 associated to the standard ℓ^1 metric on each sector $\sigma_c \subset \mathbb{R}^n$. Let us also denote by d_1 the standard metric on \mathbb{R}^n . We claim that if $\beta = [b, v] \leq_M \gamma = [c, w]$, then $d_1(\beta, \gamma) = d_1(v, w)$. First notice that the second projection $M_{\mathbb{R}} \to \mathbb{R}$ is 1–Lipschitz with respect to the metrics d_1 ; hence we have $d_1(\beta, \gamma) \geq d_1(v, w)$. By the definition of the order relation \leq_M , there exists a sequence $\beta_0 = \beta \leq_M \beta_1 \leq_M \cdots \leq_M \beta_p = \gamma$ such that, for each $0 \leq i \leq p - 1$, the points β_i, β_{i+1} lie in a common sector σ_{c_i} . Let us write representatives $\beta_i = [b_i, v_i]$, with $b_i \in L$ and $v_i \in \sigma$, for $0 \leq i \leq p$. We deduce that, for each $0 \leq i \leq p - 1$ we have $d_1(\beta_i, \beta_{i+1}) = d_1(v_i, v_{i+1})$; hence

$$d_1(\beta, \gamma) \leq \sum_{i=0}^{p-1} d_1(\beta_i, \beta_{i+1}) \leq \sum_{i=0}^{p-1} d_1(v_i, v_{i+1}) = d_1(v, w).$$

So we have proved that if $\beta = [b, v] \leq_M \gamma = [c, w]$, then $d_1(\beta, \gamma) = d_1(v, w)$.

Algebraic & Geometric Topology, Volume 24 (2024)

We now return to the increasing sequence $(\alpha_k)_{k\geq 0}$ in $M_{\mathbb{R}}$, bounded above by some $\alpha \in M_{\mathbb{R}}$. For each $k \in \mathbb{N}$, let us consider a representative $\alpha_k = [a_k, u_k]$ of α_k , and a representative $\alpha = [a, u]$ of α . Since the sequence $(\alpha_k)_{k\geq 0}$ is increasing in $M_{\mathbb{R}}$, we deduce that the sequence $(u_k)_{k\geq 0}$ is increasing in \mathbb{R}^n , and bounded above by u. Since increasing sequences in \mathbb{R}^n are geodesics for the metric d_1 , we deduce that, for each $0 \leq j \leq k$, we have $d_1(u_j, u_k) + d_1(u_k, u) = d_1(u_j, u)$.

Then, for each $0 \le j \le k$, according to the claim about the metric d_1 , we have $d_1(\alpha_j, \alpha_k) + d_1(\alpha_k, \alpha) = d_1(\alpha_j, \alpha)$. In particular, the sequence $(\alpha_k)_{k\ge 0}$ is a Cauchy sequence in $(M_{\mathbb{R}}, d_1)$. Since $M_{\mathbb{R}}$ has finitely many shapes, the proof of [Bridson and Haefliger 1999, Theorem 7.13] applies to show that the metric space $(M_{\mathbb{R}}, d_1)$ is complete; hence the sequence $(\alpha_k)_{k\ge 0}$ converges in $M_{\mathbb{R}}$. Equivalently, we can apply [Bridson and Haefliger 1999, Theorem 7.13] to the metric space $M_{\mathbb{R}}$ endowed with the length metric d_2 associated to the standard ℓ^2 metric on each sector $\sigma_c \subset \mathbb{R}^n$. Since d_1 and d_2 are bi-Lipschitz, this also implies that the metric space $(M_{\mathbb{R}}, d_1)$ is complete.

So every upper bounded subset of $M_{\mathbb{R}}$ has a join.

We have proved that $M_{\mathbb{R}}$ is a lattice such that each upper bounded subset has a join. There is an increasing action of \mathbb{R} on $M_{\mathbb{R}}$ satisfying the assumptions of Theorem 2.1 according to Lemma 3.7. As a consequence, we deduce that the metric space $(M_{\mathbb{R}}, d)$ is injective, with respect to the metric,

for all
$$x, y \in M_{\mathbb{R}}$$
, $d(x, y) = \inf\{t \ge 0 \mid (-t) \cdot x \le y \le t \cdot x\}$.

Note that the metric *d* is geodesic, and it restricts on each $\sigma_c \subset M_{\mathbb{R}}$, for $c \in C(L)$, to the natural ℓ^{∞} metric on $\sigma_c \subset \mathbb{R}^n$. Therefore *d* coincides with the length metric $d_{\mathbb{R}}$.

So we conclude that $(M_{\mathbb{R}}, d_{\mathbb{R}})$ is injective.

We saw in the Introduction that the existence of a bicombing may be extremely useful, notably in the case of Deligne complexes of Artin groups. Let us recall that a *geodesic bicombing* on a metric space X is a map $\sigma: X \times X \times [0, 1] \rightarrow X$ such that, for all $x, y \in X$, the map $t \in [0, 1] \mapsto \sigma(x, y, t)$ is a constant-speed geodesic from x to y.

The bicombing σ is called

- *reversible* if for all $x, y \in X$, for all $t \in [0, 1]$, $\sigma(x, y, t) = \sigma(y, x, 1-t)$,
- consistent if for all $x, y \in X$, for all $r, s, t \in [0, 1]$, $\sigma(\sigma(x, y, r), \sigma(x, y, s), t) = \sigma(x, y, (1-t)r + ts)$,
- *conical* if for all $x, x', y, y' \in X$, for all $t \in [0, 1]$, $d(\sigma(x, y, t), \sigma(x', y', t)) \leq (1-t)d(x, x') + td(y, y')$,
- *convex* if for all $x, x', y, y' \in X$, the map $t \in [0, 1] \mapsto d(\sigma(x, y, t), \sigma(x', y', t))$ is convex.

Note that any consistent, conical bicombing is convex.

Theorem 3.9 The metric space $(M_{\mathbb{R}}, d_{\mathbb{R}})$ has a unique convex, consistent, reversible geodesic bicombing.

Proof Let us write $X = M_{\mathbb{R}}$ for simplicity. Given any $x, y \in X$, let us define

$$D(x, y) = \inf\{t \in \mathbb{R} \mid x \le t \cdot y\} \in \mathbb{R}.$$

Since the action of \mathbb{R} on $M_{\mathbb{R}}$ is continuous, this infimum is attained; hence $x \leq D(x, y) \cdot y$. Note that this quantity is not symmetric with respect to x and y, and we have,

for all
$$x, y \in X$$
, $d(x, y) = \max(|D(x, y)|, |D(y, x)|)$.

We will start by defining a conical bicombing σ on X with a nice property which we call lower consistency. We then show that this is sufficient to bypass the use of properness of X in [Basso 2024, Theorem 1.4] applied to σ .

Fix $x, y \in X$, $t \in [0, 1]$, and let D = d(x, y). For any $a \in X$, since $-D \cdot x \leq y \leq D \cdot x$, we have $|D(x, a) - D(y, a)| \leq D$, and so

$$-tD \cdot x \leq (-tD(x,a) + tD(y,a)) \cdot x$$
$$\leq (-tD(x,a) + tD(y,a)) \cdot (D(x,a) \cdot a)$$
$$\leq ((1-t)D(x,a) + tD(y,a)) \cdot a.$$

Since every nonempty subset of X with a lower bound has a meet, we may thus define

$$\sigma(x, y, t) = \bigwedge_{a \in X} \left((1 - t)D(x, a) + tD(y, a) \right) \cdot a.$$

We will first prove that it defines a geodesic bicombing, ie that $d(x, \sigma(x, y, t)) = tD$. We have proved that, for every $a \in X$, we have $-tD \cdot x \leq ((1-t)D(x, a) + tD(y, a)) \cdot a$; hence $-tD \cdot x \leq \sigma(x, y, t)$. Conversely, when x = a, we know that

$$\sigma(x, y, t) \leq ((1-t)D(x, x) + tD(y, x)) \cdot x \leq tD(y, x) \cdot x \leq tD \cdot x.$$

We conclude that $d(x, \sigma(x, y, t)) \leq tD$. By symmetry, we also have $d(\sigma(x, y, t), y) \leq (1-t)D$. Since d(x, y) = D, we conclude that $d(x, \sigma(x, y, t)) = tD$. So σ is a geodesic bicombing. It is clear that σ is reversible.

We will now prove that σ is conical. Fix $x, y, z \in X$, and $t \in [0, 1]$. For any $a \in X$, we have $|D(y, a) - D(z, a)| \le d(y, z)$; hence $D(y, a) \le D(z, a) + d(y, z)$. We deduce that

$$\sigma(x, y, t) = \bigwedge_{a \in X} \left((1 - t)D(x, a) + tD(y, a) \right) \cdot a$$

$$\leq \bigwedge_{a \in X} \left((1 - t)D(x, a) + tD(z, a) + td(y, z) \right) \cdot a \leq td(y, z) \cdot \sigma(x, z, t).$$

By symmetry, we also have $\sigma(x, z, t) \leq t d(y, z) \cdot \sigma(x, y, t)$; hence $d(\sigma(x, y, t), \sigma(x, z, t)) \leq t d(y, z)$. So the bicombing σ is conical.

We will now prove that σ is what we will call *lower consistent*, which is one part of the inequality of the consistency equality. For each $x, y \in X$ and $s, t \in [0, 1]$, we will prove that

$$\sigma(x,\sigma(x,y,t),s) \leq \sigma(x,y,st).$$

Algebraic & Geometric Topology, Volume 24 (2024)

Let us set $z = \sigma(x, y, t)$ and $w = \sigma(x, \sigma(x, y, t), s)$: we want to prove that $w \le \sigma(x, y, st)$. For each $a \in X$, we have

$$D(w,a) \leq (1-s)D(x,a) + sD(z,a)$$

$$\leq (1-s)D(x,a) + s((1-t)D(x,a) + tD(y,a))$$

$$\leq (1-st)D(x,a) + stD(y,a).$$

Hence we deduce that $w \leq D(w, a) \cdot a \leq (1 - st)D(x, a) \cdot a + stD(y, a) \cdot a$. Since this holds for any $a \in X$, we conclude that $w \leq \sigma(x, y, st)$. Hence σ is lower consistent.

According to [Basso 2024, Lemma 5.2], given any $x, y \in X$ and $n \ge 1$, there exist unique elements $\sigma_{xy}(n, i)$, for $0 \le i \le n$, such that $\sigma_{xy}(n, 0) = x$, $\sigma_{xy}(n, n) = y$, and,

for all
$$1 \le i \le n-1$$
, $\sigma_{xy}(n,i) = \sigma(\sigma_{xy}(n,i-1), \sigma_{xy}(n,i+1), \frac{1}{2})$.

Note that even though [Basso 2024, Lemma 5.2] is stated for a proper metric space, the uniqueness part only requires that σ is a conical bicombing. And also the remark after the proof tells us that the only property needed for the existence is that the space is complete. We will actually give a proof below for the existence part.

Fix $n \ge 1$, we will prove that the elements $\sigma_{xy}(n, i)$ exist. For each $0 \le i \le n$, let us define $x_i^0 = \sigma(x, y, \frac{i}{n})$. For each $k \in \mathbb{N}$, let us define $x_0^k = x$ and $x_n^k = y$. For each $k \in \mathbb{N}$ and $1 \le i \le n-1$, let us define inductively $x_i^{k+1} = \sigma(x_{i-1}^k, x_{i+1}^k, \frac{1}{2})$. Since σ is lower consistent, we see by induction that, for each $0 \le i \le n$, the sequence $(x_i^k)_{k \in \mathbb{N}}$ is nonincreasing in *X*. Moreover, since each x_i^k lies on a geodesic from *x* to *y*, we have $x_i^k \ge (-d(x, y)) \cdot x$ for every $k \in \mathbb{N}$ and $0 \le i \le n$.

Hence for each $0 \le i \le n$, since the sequence $(x_i^k)_{k \in \mathbb{N}}$ has a lower bound, we may define its meet $\sigma_{xy}(n,i) = \bigwedge_{k\ge 0} x_i^k$. In fact, the sequence $(x_i^k)_{k\ge 0}$ actually converges to $\sigma_{xy}(n,i)$. By the continuity of σ , we deduce that the elements $(\sigma_{xy}(n,i))_{0\le i\le n}$ satisfy the required property.

Note that [Basso 2024, Theorem 1.4] is stated for a proper metric space, but the properness assumption is used in precisely two arguments: first in [Basso 2024, Lemma 5.2] to prove the existence of the elements $\sigma_{xy}(n, i)$, which we obtained using specific properties of X.

Properness of X is used again, although not explicitly stated, in the proof of [Basso 2024, Theorem 1.4] to ensure the pointwise convergence of a sequence of bicombings with respect to some ultrafilter. Instead of using ultrafilters to ensure convergence, we will rather use the lower consistency of the bicombing.

For each $n \ge 1$, [Basso 2024, Lemma 5.2] states that the function $\sigma^{(n)}: X \times X \times [0, 1] \to X$, defined by

$$\sigma^{(n)}\left(x, y, (1-\lambda)\frac{i}{n} + \lambda\frac{i+1}{n}\right) = \sigma(\sigma_{xy}(n, i), \sigma_{xy}(n, i+1), \lambda)$$

for all $x, y \in X$, $\lambda \in [0, 1]$ and $0 \le i \le n - 1$, is a conical bicombing.

First note that, since σ is lower consistent, and by the uniqueness of the points $\sigma_{xy}(n, i)$, we have for all $x, y \in X, n \ge 1, 0 \le i \le n-1$ and $p \ge 1$ that $\sigma_{xy}(np, ip) \le \sigma_{xy}(n, i)$. For each $x, y \in X$ and $t \in [0, 1]$, let us define

$$\gamma(x, y, t) = \bigwedge_{n \ge 1} \sigma_{xy}(n, \lceil tn \rceil).$$

According to the previous property, we deduce that

$$\gamma(x, y, t) = \lim_{n \to +\infty} \sigma_{xy}(n!, \lceil tn! \rceil).$$

Since each $\sigma^{(n)}$ is a conical bicombing, one also deduces that γ is conical.

We will prove that γ is lower consistent: Let $x, y \in X$ and $s, t \in [0, 1]$. We have

$$\gamma(x, \gamma(x, y, t), s) \leq \lim_{n \to +\infty} \gamma(x, \sigma_{xy}(n!, \lceil tn! \rceil), s)$$

$$\leq \lim_{n, m \to +\infty} \sigma_{xy}(n! m!, \lceil s \lceil tn! \rceil m! \rceil)$$

$$\leq \lim_{n, m \to +\infty} \sigma_{xy}(n! m!, \lceil stn! m! \rceil) \leq \gamma(x, y, st)$$

$$(\sigma_{xy}(n! m!, \lceil s \lceil tn! \rceil m! \rceil), \sigma_{xy}(n! m!, \lceil stn! m! \rceil)) \leq \frac{\lceil sm! \rceil}{n! m!} d(x, y) \to 0$$

as

d

as
$$n \to +\infty$$
. So we deduce that γ is a reversible, conical, lower consistent geodesic bicombing such that $\gamma \leq \sigma$.

As a consequence, if we start with a reversible, conical, lower consistent geodesic bicombing σ' which is minimal, we have $\gamma' = \sigma'$; hence σ' satisfies the following consistency property: for all $x, y \in X$, for all $s, t \in [0, 1]$, $\sigma'(x, \sigma'(x, y, t), s) = \sigma'(x, y, st)$. Since σ' is reversible, we deduce that σ' is actually consistent. Since σ' is conical, it is also convex.

According to [Descombes and Lang 2015, Theorem 1.2], since X has finite combinatorial dimension, we conclude that σ' is the unique convex consistent reversible geodesic bicombing of X.

Theorem 3.10 Let *L* denote a bounded, graded lattice. The orthoscheme realization |L| of *L*, endowed with the piecewise ℓ^{∞} metric, is injective. Moreover, |L| has a unique convex reversible consistent geodesic bicombing.

Proof Consider the affine version $M = M_{\mathbb{R}}$ of L over \mathbb{R} . For some maximal chain $c \in C(L)$, consider the elements $0_M = [(0, \ldots, 0), c] \in M$, $\mu_M = [(\frac{1}{2}, \ldots, \frac{1}{2}), c] \in M$ and $1_M = [(1, \ldots, 1), c] \in M$: note that 0_M , μ_M and 1_M do not depend on c.

Note that the interval $I(0_M, 1_M)$ coincides with the ball $B(\mu_M, \frac{1}{2})$ in M for the metric $d_{\mathbb{R}}$. According to Theorem 3.8, $(M, d_{\mathbb{R}})$ is injective. So the ball $B(\mu_M, \frac{1}{2}) = I(0_M, 1_M)$ is injective.

We remark that the interval $I(0_M, 1_M)$ of M, endowed with the metric $d_{\mathbb{R}}$, is isometric to the orthoscheme realization of L, endowed with the piecewise ℓ^{∞} metric. Indeed, for each maximal chain $c \in C(L)$,

Algebraic & Geometric Topology, Volume 24 (2024)

notice that $\sigma_C \cap I(0_M, 1_M)$ identifies with the standard orthoscheme $\{x \in \mathbb{R}^n \mid 0 \le x_1 \le \ldots \le x_n \le 1\}$, where *n* denotes the rank of *L*. We conclude that the orthoscheme realization of *L* is injective.

According to Theorem 3.9, we know that $M_{\mathbb{R}}$ has a unique convex reversible consistent geodesic bicombing σ . Since the ball $|L| = B(\mu_M, \frac{1}{2})$ is stable under σ , we deduce that |L| has a convex reversible consistent geodesic bicombing. Since |L| has combinatorial dimension at most n, according to [Descombes and Lang 2015, Theorem 1.2], we deduce that σ is the only convex bicombing on |L|. \Box

Note that knowing when the orthoscheme complex of a lattice, endowed with the piecewise Euclidean metric, is CAT(0) is a very subtle question (see the Introduction). We can prove the following.

Theorem 3.11 Let *L* denote a bounded, graded poset, let |L| denote the geometric realization of *L*, and let d_p denote the ℓ^p orthoscheme metric on |L| for $p \in \{2, \infty\}$. If $(|L|, d_2)$ is CAT(0), then $(|L|, d_{\infty})$ is injective. The converse is false.

Proof Assume that *L* is a bounded, graded poset such that $(|L|, d_{\infty})$ is not injective. According to Theorem 3.10, we deduce that *L* is not a lattice. According to Proposition 1.5, there exists a bowtie in *L*: consider $x_1, x_2, x_3, x_4 \in L$ such that x_1 and x_3 are maximal elements inferior to both x_2 and x_4 , and such that x_2 and x_4 are minimal elements superior to both x_1 and x_3 . In the link $L_{[0,1]}$ of the diagonal edge [0, 1] in $(|L|, d_2)$, consider the piecewise geodesic loop ℓ going through x_1, x_2, x_3, x_4, x_1 .

According to [Brady and McCammond 2010, Proposition 4.8], each geodesic segment $[x_i, x_{i+1}]$ has length smaller than $\frac{\pi}{2}$; hence ℓ has length smaller than 2π . On the other hand, according to Lemma 7.2 of that work the loop ℓ is locally geodesic in $L_{[0,1]}$. Hence $L_{[0,1]}$ is not CAT(1), and $(|L|, d_2)$ is not CAT(0).

We will now prove that the converse does not hold. Consider the Coxeter symmetric group $W = \mathfrak{S}_4$, with standard generators $S = \{s_1, s_2, s_3\}$ and standard Garside longest element $\Delta = s_1 s_2 s_3 s_1 s_2 s_1$. Let L denote the poset W, with order relation "being a left prefix for a shortest representative in S". Then L is a lattice, and so $(|L|, d_{\infty})$ is injective according to Theorem 3.10.

On the other hand, consider the piecewise geodesic loop ℓ in the link $L_{[0,1]}$ of the diagonal edge [0, 1] in $(|L|, d_2)$ going through the vertices s_1 , $s_1s_2s_1$, s_2 , $s_2s_3s_2$, s_3 , s_3s_1 , s_1 . According to [Brady and McCammond 2010, Proposition 4.8], its length is

$$2 \arccos\left(\sqrt{\frac{1}{2}\frac{4}{5}}\right) + 4 \arccos\left(\sqrt{\frac{1}{3}\frac{3}{5}}\right) \simeq 0.987(2\pi) < 2\pi.$$

Since $L_{[0,1]}$ has the homotopy type of a circle, ℓ is not nullhomotopic in $L_{[0,1]}$. So $L_{[0,1]}$ is not CAT(1), and $(|L|, d_2)$ is not CAT(0).

We also deduce an immediate consequence concerning the Garside complex of a Garside group. Fix a Garside group (G, S, δ) . Let X denote the *Garside complex* of G, ie the simplicial complex with vertex set G, and with simplices corresponding to chains $g_1 <_L g_2 <_L \cdots <_L g_n$ such that $g_n \leq_L g_1 \delta$. Since simplices of X have a total order on their vertices, X is a simplicial complex with ordered simplices, so we may endow X with the piecewise ℓ^{∞} orthoscheme metric.

Corollary 3.12 The Garside complex of any Garside group, endowed with the piecewise ℓ^{∞} orthoscheme metric, is injective.

This applies, in particular, to the dual braid complex studied by Brady and McCammond [2010]. This complex, endowed with the piecewise orthoscheme Euclidean metric, is conjectured to be CAT(0), but it is only known for a very small number of strands (see [Brady and McCammond 2010; Haettel et al. 2016; Jeong 2023]). On the other hand, if we endow the dual braid complex with the piecewise orthoscheme ℓ^{∞} metric, we see that it is injective.

4 Application to Euclidean buildings and the Deligne complex of type $\tilde{A_n}$

Let us consider the Euclidean Coxeter group $W \simeq \mathfrak{S}_n \ltimes \mathbb{Z}^{n-1}$ of type \tilde{A}_{n-1} . Its Coxeter complex may be identified with

$$\Sigma = \{ x \in \mathbb{R}^n \mid x_1 + \dots + x_n = 0 \}.$$

Up to homothety, we may choose the following affine hyperplanes to define Σ :

$$\{x_i - x_j = k \mid 1 \leq i \neq j \leq n, k \in \mathbb{Z}\},\$$

so that maximal simplices of Σ identify with the W-orbit of

$$K = \{ x \in \Sigma \mid x_1 \leqslant x_2 \leqslant \ldots \leqslant x_n \leqslant x_1 + 1 \}.$$

The vertex set of Σ identifies with $\left(\frac{1}{n}\mathbb{Z}\right)^n \cap \Sigma$. Since the simplex *K* is a strict fundamental domain for the action of *W* on Σ , one may define a *W*-invariant type function τ on the vertex set of Σ :

$$\tau: \Sigma^{(0)} \to \mathbb{Z}/n\mathbb{Z}, \quad x \in W \cdot v_i \mapsto i,$$

where $v_i = \left(\frac{n-i}{n}, \dots, \frac{n-i}{n}, -\frac{i}{n}, \dots, -\frac{i}{n}\right)$ is the vertex of *K* whose first *i* coordinates equal $\frac{n-i}{n}$ and the n-i last coordinates equal $-\frac{i}{n}$.

This type of function is such that adjacent vertices have distinct types.

Let us define the extended Coxeter complex

$$\widehat{\Sigma} = \Sigma \times \mathbb{R} = \mathbb{R}^n,$$

where the action of the standard generators w_1, \ldots, w_n of W on $\hat{\Sigma}$ is given by

for all
$$1 \le i \le n-1$$
, $w_i \cdot (x_1, \dots, x_i, x_{i+1}, \dots, x_n) = (x_1, \dots, x_{i+1}, x_i, \dots, x_n)$
and $w_n \cdot (x_1, \dots, x_n) = (x_n - 1, x_2, \dots, x_{n-1}, x_1 + 1)$.

Also note that $\hat{\Sigma}$ has a natural simplicial complex structure, with vertex set \mathbb{Z}^n , with maximal simplices corresponding to the *W*-orbits of the simplices

$$\{k \leq x_i \leq x_{i+1} \leq x_{i+2} \leq \ldots \leq x_{i+n-1} \leq k+1 \mid k \in \mathbb{Z}, i \in \mathbb{Z}/n\mathbb{Z}\},\$$

where indices in \mathbb{R}^n are considered modulo *n*. Note that the action of the Coxeter group *W* preserves τ .

Algebraic & Geometric Topology, Volume 24 (2024)

A fundamental domain for the action of W on Σ is the simplex

$$K = \{ x \in \Sigma \mid x_1 \leq x_2 \leq \ldots \leq x_n \leq x_1 + 1 \},\$$

and a fundamental domain for the action of W on $\hat{\Sigma}$ is the column

$$\widehat{K} = \{ x \in \widehat{\Sigma} \mid x_1 \leq x_2 \leq \cdots \leq x_n \leq x_1 + 1 \}.$$

Note that such columns have been studied by Brady and McCammond [2010], and by Dougherty, McCammond and Witzel [Dougherty et al. 2020].

We will endow $\hat{\Sigma}$ with the standard ℓ^{∞} metric from \mathbb{R}^n .

Let us consider a simplicial complex X such that

- either X is a Euclidean building of type \tilde{A}_{n-1} ,
- or X is the Deligne complex of the Euclidean Artin group A of type \tilde{A}_{n-1} , with a coarser simplicial structure.

We will define the *extended* version of X, denoted by \hat{X} . It is a simplicial complex whose geometric realization is homeomorphic to $X \times \mathbb{R}$.

Before giving the precise definition of \hat{X} , let us first recall briefly the definition of the Deligne complex Δ of the Euclidean Artin group A of type \tilde{A}_{n-1} (see also Section 1.2). Let $S \simeq \mathbb{Z}/n\mathbb{Z}$ denote the standard generating set of A. Consider the set

$$\{gA_T \mid g \in A, T \subsetneq S\},\$$

endowed with the following partial order: $gA_T \leq g'A_{T'}$ if $gA_T \subset g'A_{T'}$. Then the Deligne complex Δ of A is the geometric realization of this poset. We will define a coarser simplicial structure X on Δ . Note that, for any minimal vertex $gA_{\emptyset} \in \Delta$ (where $g \in A$), the 1–neighborhood of gA_{\emptyset} in Δ is

$$\{gA_T \mid T \subsetneq S\}$$

which is precisely the barycentric subdivision of the simplex with vertex set

$$\{gA_T \mid T \subsetneq S, |S \setminus T| = 1\}$$

consisting of (the g-translates of) all maximal proper standard parabolic subgroups of A.

Therefore we may define the simplicial complex X with vertex set $\{gA_T \mid g \in A, T \subsetneq S, |S \setminus T| = 1\}$, and such that $g_1A_{T_1}, \ldots, g_kA_{T_k}$ span a simplex in X if and only if $\bigcap_{i=1}^k g_iA_{T_i} \neq \emptyset$. Then Δ identifies with the barycentric subdivision of X, and also the geometric realizations of Δ and X are homeomorphic.

Note that, in both cases (Euclidean building or Deligne complex), there is a well-defined type function $\tau: X^{(0)} \to \mathbb{Z}/n\mathbb{Z}$ such that adjacent vertices have different types.

• In the case X is a Euclidean building of type \tilde{A}_{n-1} , each apartment is identified with the Coxeter complex Σ ; we may define the type of a vertex v of X to be its type in any apartment containing v. Since the Coxeter group W preserves the type, this definition does not depend on the choice of apartment.

• In the case X is the Deligne complex of type \tilde{A}_{n-1} , one may either use the projection onto the Coxeter complex, or use a direct definition: if gA_T is a vertex of X, with $g \in A$ and $T = S \setminus \{i\}$, its type is *i*.

The complex X may also be defined as a particular gluing of copies of the Coxeter complex Σ . Roughly speaking, \hat{X} will be the same gluing of copies of $\hat{\Sigma}$.

More precisely, let $\hat{X} = X \times \mathbb{R}$, and we will define a simplicial complex structure on \hat{X} . The vertex set of \hat{X} is $\{(x, i) \in X^{(0)} \times \mathbb{Z} \mid \tau(x) = i + n\mathbb{Z}\}$, where $X^{(0)}$ denotes the vertex set of X. Using the type function τ , remark that vertices of maximal simplices of X have a well-defined cyclic ordering in $\mathbb{Z}/n\mathbb{Z}$. The maximal simplices of \hat{X} are

$$\{(x_i, kn+i), (x_{i+1}, kn+i+1), \dots, (x_n, kn+n), (x_1, kn+n+1), (x_2, kn+n+2), \dots, (x_i, kn+n+i)\}$$

for any $k \in \mathbb{Z}$, any $1 \le i \le n$ and any maximal simplex (x_1, x_2, \ldots, x_n) of X with for all $1 \le i \le n$, $\tau(x_i) = i$. We endow each such simplex with the ℓ^{∞} orthoscheme metric for the given ordering. Endow \hat{X} with the associated length metric.

Note that the translation action on the \mathbb{R} factor of $\hat{X} = X \times \mathbb{R}$ defines an isometric action denoted by θ . Also note that θ does not preserve the simplicial structure of \hat{X} , but only its restriction to the subgroup $n\mathbb{Z}$ of \mathbb{R} .

If X is a Euclidean building of type \tilde{A}_{n-1} , then \hat{X} is called a Euclidean building of extended type \tilde{A}_{n-1} (see [Bruhat and Tits 1972], and also [Haettel 2022a]).

If X is the Deligne complex of the Euclidean Artin group A of type \tilde{A}_{n-1} , we will give another description of \hat{X} . Recall that the classical Deligne complex X may be defined as

$$X = (A \times K) / \sim,$$

where $(g, x) \sim (g', x')$ if x = x' and, if the stabilizer of x in Σ equals W_T for some $T \subsetneq S$, then $g^{-1}g' \in A_T$. Then the extended Deligne complex may also be defined similarly as

$$\widehat{X} = (A \times \widehat{K}) / \sim,$$

where $(g, x) \sim (g', x')$ if x = x' and, if the stabilizer of x in $\hat{\Sigma}$ equals W_T for some $T \subsetneq S$, then $g^{-1}g' \in A_T$.

Fix any vertex $x \in X$, and let $L_{x,0}$ denote the set of vertices of X adjacent to x. Without loss of generality, we may assume that x has type $\tau(x) = 0$. Note that vertices in $L_{x,0}$ have a type in $\mathbb{Z}/n\mathbb{Z}\setminus\{\tau(x)\} = \{1, \ldots, n\}$, which is an interval. Since maximal simplices of X have a natural cyclic ordering in $\mathbb{Z}/n\mathbb{Z}$, there is a natural induced order on $L_{x,0}$ that is consistent with the type function $\tau: L_{x,0} \to \{1, \ldots, n\}$. Consider the poset $L_x = L_{x,0} \cup \{0, 1\}$, where 0 and 1 are defined to be the minimum and the maximum of L_x respectively.

Proposition 4.1 Consider any $p \in \hat{X}$ whose projection p_X onto X is contained in the open star of the vertex x. Then \hat{X} is locally isometric at p to a neighborhood of a point in the ℓ^{∞} orthoscheme realization of the poset L_x .

Algebraic & Geometric Topology, Volume 24 (2024)

Proof Remember that there is a diagonal, isometric action θ of \mathbb{R} on \hat{X} , whose quotient is X. So, up to isometry, we may assume that p lies in the open star of the diagonal edge joining the vertices (x, 0) and (x, n) of \hat{X} . Note that each simplex of \hat{X} containing p corresponds to a chain in L_x containing (x, 0) (identified with $0 \in L_x$) and (x, n) (identified with $1 \in L_x$). As a consequence, a neighborhood of p in \hat{X} is isometric to a neighborhood of a point in the orthoscheme realization of L_x .

We now prove that, in the case of a Euclidean building, the poset L_x is a lattice.

Proposition 4.2 Let L_0 denote the vertex set of a (possibly nonthick) spherical building of type A_{n-1} for some $n \ge 1$, and let $L = L_0 \cup \{0, 1\}$. In other words, L is the poset of linear subspaces of a projective space of dimension n - 1. Then L is a bounded, graded lattice of rank n - 1.

Proof The lattice property is obvious: if S, S' are linear subspaces of a projective space X, then $S \wedge S' = S \cap S'$ and $S \vee S'$ is the linear subspace spanned by $S \cup S'$.

As a consequence, if X is a Euclidean building of type \tilde{A}_{n-1} , for any vertex $x \in X$, the poset L_x is a bounded graded lattice.

We now turn to the case of the Deligne complex of type \tilde{A}_{n-1} . We defer the proof of the following result to Section 5.

Theorem 4.3 (Crisp–McCammond) Let X denote the Deligne complex of type \tilde{A}_{n-1} . For any vertex $x \in X$, the poset L_x is a bounded, graded lattice.

We may now deduce the main result.

Theorem 4.4 Any Euclidean building of extended type \tilde{A}_{n-1} , or the extended Deligne complex of type \tilde{A}_{n-1} , endowed with the piecewise ℓ^{∞} metric, is injective.

Proof According to Proposition 4.1 and Theorem 4.3, we know that \hat{X} is locally isometric to the ℓ^{∞} orthoscheme complex of a bounded graded lattice. According to Theorem 3.10, we deduce that \hat{X} is uniformly locally injective.

If X is a Euclidean building, X and \hat{X} are contractible, and in particular simply connected. If X is a Deligne complex, according to Theorem 1.6, \hat{X} is simply connected.

According to Theorem 1.11, \hat{X} is complete. We deduce with Theorem 1.14 that \hat{X} is injective.

We can also deduce that the thickening is a Helly graph. Note that this thickening, described more precisely in Theorem 1.15, corresponds to a coarser cell structure on the building or the Deligne complex.

Corollary 4.5 The thickening of the vertex set of any Euclidean building of extended type \tilde{A}_{n-1} , or the extended Deligne complex of type \tilde{A}_{n-1} , is a Helly graph.

Proof Let X denote either a Euclidean building of extended type \tilde{A}_{n-1} or the extended Deligne complex of type \tilde{A}_{n-1} . We will see that X satisfies the assumptions of Theorem 1.15.

Since simplices of X have a well-defined order, we see that X is a simplicial complex with ordered simplices. For each maximal simplex σ of X, the minimal and maximal vertices of σ form a maximal edge in X. Therefore, X has maximal edges as in Definition 1.9. According to Theorem 1.15, we deduce that the thickening of X is a Helly graph.

We will now deduce a bicombing on X, considered as the quotient of $(\hat{X}, d_{\hat{X}})$ by the diagonal action θ of \mathbb{R} on \hat{X} . Let us define the quotient metric d_X on X:

for all
$$x, y \in X$$
, $d_X(x, y) = \inf_{t \in \mathbb{R}} d_{\widehat{X}}(x, \theta(t) \cdot y)$.

Theorem 4.6 Any Euclidean building X of type \tilde{A}_{n-1} , or the Deligne complex X of type \tilde{A}_{n-1} , has a metric d_X that admits a convex, consistent, reversible geodesic bicombing σ . Moreover, d_X and σ are invariant under the group of type-preserving automorphisms of X.

Proof We will prove it locally, ie for the star Y of a vertex v of X. According to Theorem 3.10, there exists a unique convex, consistent, reversible bicombing $\hat{\sigma}$ on \hat{Y} . For each $x, y \in Y$, choose lifts $\hat{x}, \hat{y} \in \hat{Y}$ such that $d_{\hat{X}}(\hat{x}, \hat{y}) = d_X(x, y)$. For each $t \in [0, 1]$, let us define $\sigma(x, y, t) = \theta(\mathbb{R}) \cdot \hat{\sigma}(\hat{x}, \hat{y}, t) \in Y$.

We will first see that σ is well-defined: indeed fix $x, y \in Y$, and consider two pairs of lifts $\hat{x}, \hat{x}', \hat{y}, \hat{y}' \in \hat{Y}$ such that $d_{\hat{X}}(\hat{x}, \hat{y}) = d_{\hat{X}}(\hat{x}', \hat{y}') = d_X(x, y)$. Note that, up to the action of θ , we may assume that $\hat{x} = \hat{x}'$. If $\hat{y} \neq \hat{y}'$, let $a \in \mathbb{R} \setminus \{0\}$ such that $\hat{y}' = \theta(a) \cdot \hat{y}$. We then have $d_{\hat{X}}(\hat{x}, \theta(\frac{a}{2}) \cdot \hat{y}) < d(x, y)$, which is a contradiction. So $\hat{y} = \hat{y}'$, and σ is well-defined.

Moreover, it is easy to see that σ is a reversible, consistent, geodesic bicombing.

We will now see that σ is convex. Consider $x, x', y, y' \in Y$, and choose lifts $\hat{x}, \hat{x}', \hat{y}, \hat{y}' \in \hat{Y}$ such that $d_{\hat{X}}(\hat{x}, \hat{x}') = d_X(x, x')$ and $d_{\hat{X}}(\hat{y}, \hat{y}') = d_X(y, y')$. Fix any $t \in [0, 1]$ and $s \in \mathbb{R}$. We have

$$d_X(\sigma(x, y, t), \sigma(x', y', t)) = \inf_{s \in \mathbb{R}} d_{\widehat{X}}(\widehat{\sigma}(\widehat{x}, \widehat{y}, t), \theta(s) \cdot \widehat{\sigma}(\widehat{x}, \widehat{y}, t))$$

$$\leq d_{\widehat{X}}(\widehat{\sigma}(\widehat{x}, \widehat{y}, t), \widehat{\sigma}(\widehat{x}, \widehat{y}, t))$$

$$\leq (1 - t) d_{\widehat{X}}(\widehat{x}, \widehat{x}') + t d_{\widehat{X}}(\widehat{y}, \widehat{y}')$$

$$\leq (1 - t) d_X(x, x') + t d_X(y, y'),$$

so σ is a conical bicombing. Since σ is also consistent, it is a convex bicombing.

We have seen that (X, d_X) is locally convexly reversibly consistently bicombable. According to [Miesch 2017, Theorem 1.1], we deduce that (X, d_X) has a unique global convex reversible geodesic consistent bicombing σ that is consistent with local bicombing. Since the local bicombing only depends on the local combinatorics of X and the type, we deduce that σ is invariant under type-preserving automorphisms of X.

Algebraic & Geometric Topology, Volume 24 (2024)

Note that we strengthened this result in [Haettel 2022b], by proving that this convex geodesic bicombing is actually unique. Our result also extends to simplicial complexes which are much more general than

5 The lattice of maximal parabolic subgroups of braid groups

This whole section is unpublished work of John Crisp and Jon McCammond, copied here with their permission. The results concerning the lattice of cut-curves are contained in [Bessis 2006] about Garside structures on free groups. For consistency, we choose to follow Crisp and McCammond's presentation instead.

5.1 The lattice of cut-curves

buildings.

Let \mathbb{D}^2 denote the unit disk in \mathbb{R}^2 , and let $\{p_1, \ldots, p_n\}$ denote a set of *n* distinct points in \mathbb{D}^2 , as shown in Figure 2. We write $\mathbb{D}_* = \mathbb{D}^2 \setminus \{p_1, \ldots, p_n\}$. Let a = (0, 1) and b = (0, -1) denote the top and bottom points of the boundary $\partial \mathbb{D}^2$.

Definition 5.1 By a *cut-curve* or *curve* on \mathbb{D}_* we shall mean a smoothly embedded curve c in \mathbb{D}_* which meets $\partial \mathbb{D} = \partial \mathbb{D}^2$ precisely at its endpoints, and which separates the boundary points a and b.

We shall say that two cut-curves are isotopic if they are isotopic in \mathbb{D}_* relative to $\{a, b\}$. We denote by [c] the isotopy class of a curve c, and write C for the set of all isotopy classes of cut-curves in \mathbb{D}_* .

Observe that any cut-curve c separates \mathbb{D}_* into two regions, an upper region containing a and a lower region containing b, and induces, in particular, a partition of the points $\{p_1, \ldots, p_n\}$ into two sets. In general we shall say that the contents of the region containing a lie *above* c and the contents of the region containing b lie *below* c. For each curve c, we write deg(c) for the number of points p_i which lie below c. Clearly this number is invariant under isotopy, and so defines a degree function on C by deg([c]) = deg(c).



Figure 2: An example of cut-curve c in the punctured disk \mathbb{D}_* .

Let c_1, c_2 be two curves. We say that c_1 and c_2 are *in minimal position* with respect to one another if they do not cobound any disk regions ("bigons") in \mathbb{D}_* . This includes triangular regions against the boundary. Any such disk regions can always be removed by modifying just one of the two curves in its isotopy class without changing the other. That is to say that, for any two curves c_1, c_2 , we can always find c'_1 such that $[c'_1] = [c_1]$ and c'_1 is in minimal position with respect to c_2 .

Here is another argument about minimal position using hyperbolic geometry. Such an argument has been used in [Bessis 2006]. If $n \ge 2$, one may also endow \mathbb{D}_* with a fixed complete hyperbolic metric. Fix $p^+ = (1, 0), p^- = (-1, 0) \in \partial \mathbb{D}$ as points on each connected component of $\partial \mathbb{D} \setminus \{a, b\}$. Then, for each isotopy class [c], we may consider the unique geodesic line in \mathbb{D}_* in the isotopy class [c] with endpoints $p^+, p^- \in \partial \mathbb{D}$. Then, for any $[c_1], [c_2] \in C$, the geodesic representatives are in minimal position.

Definition 5.2 Let c_1, c_2 be two curves. We say that $[c_1] \leq [c_2]$ if c_1 is isotopic to a curve which lies below c_2 . It is easily checked that this defines a partial order on the set C of cut-curves classes.

Note that if c_1 and c_2 are in minimal position with respect to one another then $[c_1] \leq [c_2]$ if and only if c_1 lies below c_2 (in particular they are disjoint). Note also that the function deg: $C \rightarrow \{1, ..., n\}$ is a strict order-preserving map, a grading on the poset (C, \leq) .

Theorem 5.3 [Bessis 2006, Theorem 2.6] The graded poset (C, \leq) is a lattice.

Proof Let $x = [c_1]$, $y = [c_2]$ be arbitrary elements of C. Suppose that the curves c_1 and c_2 are in minimal position with respect to one another. Now consider how the union of c_1 and c_2 cut \mathbb{D}_* up into connected regions. There is a unique lowermost such region R which lies below both c_1 and c_2 (and contains the point b), and another uppermost such region R' which lies above both c_1 and c_2 (and contains the point b), and another uppermost such region R' which lies above both c_1 and c_2 (and contains the point a). Let c, resp. c', denote the curves which skirt along the part of the boundary of R, resp. R', lying in the interior of \mathbb{D}_* (ie not along $\partial \mathbb{D}$). Then we claim that $x \wedge y = [c]$ and $x \vee y = [c']$.

We check just the first of these two claims. Suppose that c_0 represents a common lower bound for x and y. Then, by a sequence of bigon-removing isotopies (or by considering the geodesic representative), we may choose the representative c_0 to be in minimal position with respect to both c_1 and c_2 . Since c_0 represents a common lower bound, c_0 lies below both c_1 and c_2 , and hence lies below the curve c. \Box

Recall that the *n*-strand braid group B_n is isomorphic to the mapping class group MCG($\mathbb{D}_*, \partial \mathbb{D}$). As a consequence, B_n acts naturally on the set C of isotopy classes of cut-curves.

Lemma 5.4 The action of B_n on C preserves the order and the degree.

Proof Let $[c] \in C$ be a curve of degree $1 \le k \le n-1$ and $g \in B_n$. Then k points among $\{p_1, \ldots, p_n\}$ lie below c. Since the action of B_n fixes the boundary of \mathbb{D}_* pointwise, we know that k points among $\{p_1, \ldots, p_n\}$ lie below g(c). Hence deg g([c]) = deg[c].



Figure 3: The base maximal chain $\alpha = (c_1, c_2, \dots, c_{n-1})$ of C.

Similarly, consider two curves $[c_1], [c_2] \in C$ such that $[c_1] \leq [c_2]$. Consider c_1, c_2 in minimal position, so that c_1 is below c_2 . For any $g \in B_n$, we have that $g(c_1)$ is below $g(c_2)$; hence $g([c_1]) \leq g([c_2])$. We conclude that the action of B_n preserves the order on C.

Note that B_n acts transitively on the set of cut-curves with fixed degree.

5.2 Cosets in braid groups

We will show how the cut-curve lattice (\mathcal{C}, \leq) can be reinterpreted in purely algebraic terms.

Definition 5.5 Recall that the braid group B_n is generated by the set of standard generators $S = \{\sigma_1, \sigma_2, \dots, \sigma_{n-1}\}$. For each $1 \le k \le n-1$, let P_k denote the maximal parabolic subgroup of B_n generated by $S \setminus \{\sigma_k\}$. Thus P_k is isomorphic to a product $B_k \times B_{n-k}$. Note also that these subgroups are distinct for distinct values of k. We define the following augmented collection of cosets in B_n :

$$\mathcal{B} = \{gP_k \mid g \in B_n, 1 \leq k \leq n-1\} \cup \{0, 1\}.$$

For $b \in \mathcal{B}$, we write $\deg(b) = k$ if $b = gP_k$ for $1 \le k \le n-1$, and $\deg(0) = 0$, $\deg(1) = n$. We define an order relation $\le_{\mathcal{B}}$ on \mathcal{B} as follows. First, define $0 \le b \le 1$ for all $b \in \mathcal{B}$. Otherwise, for $g_1, g_2 \in B_n$ and $1 \le k_1, k_2 \le n-1$, we write $g_1P_{k_1} \le g_2P_{k_2}$ if $k_1 \le k_2$ and $g_1P_{k_1} \cap g_2P_{k_2} \ne \emptyset$.

For simplicity, we shall henceforth identify B_n with the mapping class group of \mathbb{D}_* (relative to $\partial \mathbb{D}$). Let us consider the base maximal chain $\alpha = (c_1, c_2, \dots, c_{n-1})$ of \mathcal{C} depicted in Figure 3.

Definition 5.6 Define a map $\Phi: C \to B$ by setting $\Phi(c) = 0$ if deg(c) = 0, $\Phi(c) = 1$ if deg(c) = n, and otherwise

$$\Phi(c) = \{g \in B_n \mid g(\alpha) \text{ contains } c\}$$

Lemma 5.7 If $c \in \alpha$ and $1 \leq \deg(c) = k \leq n - 1$, then $\Phi(c) = P_k$.

Proof The curve *c* is a base curve such that p_1, \ldots, p_k lie below *c* and p_{k+1}, \ldots, p_n lie above *c*. Let R^- denote the connected component of $\mathbb{D}^* \setminus c$ below *c*, and R^+ denote the connected component of $\mathbb{D}^* \setminus c$ above *c*. The stabilizer of *c* in the mapping class group $B_n = \text{MCG}(\mathbb{D}_*, \partial \mathbb{D})$ is the direct product of $\text{MCG}(R^-, \partial R^-) = \langle \sigma_1, \ldots, \sigma_{k-1} \rangle$ and $\text{MCG}(R^+, \partial R^+) = \langle \sigma_{k+1}, \ldots, \sigma_{n-1} \rangle$. Hence the stabilizer of *c* in B_n equals P_k .

Lemma 5.8 The map $\Phi: \mathcal{C} \to \mathcal{B}$ is well-defined, surjective, and respects degrees.

Proof Fix $c \in C$ with $1 \leq \deg(c) = k \leq n-1$. Fix some $g \in B_n$ such that the curve *c* lies in $g(\alpha)$. Let $c_0 = g^{-1}(c)$, the degree-*k* curve in α . For any $h \in B_n$, we have

$$h \in \Phi(c) \iff c \in h(\alpha) \iff g(c_0) = h(c_0).$$

According to Lemma 5.7, this is equivalent to $gP_k = hP_k$. Hence $\Phi(c) = gP_k$ is well-defined. Moreover it is clear that Φ is surjective, and that for all $c \in C$ we have $\deg(\Phi(c)) = \deg(c)$.

Lemma 5.9 The map Φ is injective, so it is a bijection between C and B.

Proof Let $c_1, c_2 \in C$ such that $\Phi(c_1) = \Phi(c_2)$. We may suppose that c_1 lies in $g_1(\alpha)$ and c_2 in $g_2(\alpha)$ for some $g_1, g_2 \in B_n$. According to Lemma 5.8, we deduce that c_1 and c_2 have the same degree $0 \leq k \leq n$. Since C and B have unique elements of degree 0 and n, we may restrict to the case $1 \leq k \leq n$.

Since $\Phi(c_1) = \Phi(c_2)$, we deduce that $g_1 P_k = g_2 P_k$, so $g_1^{-1} g_2 \in P_k$. Let $h \in P_k$ such that $g_2 = g_1 h$. Let c_0 denote the degree-k curve of α . Then $c_1 = g_1(c_0)$ and $c_2 = g_2(c_0)$. Since $h \in P_k$, h fixes c_0 so $c_1 = c_2$.

Theorem 5.10 (Crisp–McCammond) The bijection $\Phi: C \to B$ is an order isomorphism between (C, \leq) and (B, \leq_B) . As a consequence, (B, \leq_B) is a lattice.

Proof To prove the theorem, we need to show that, for $c_1, c_2 \in C$, we have $c_1 \leq c_2$ if and only if $\Phi(c_1) \leq_{\mathcal{B}} \Phi(c_2)$. Write $k_i = \deg(c_i)$ for i = 1, 2, and suppose, without loss of generality, that $0 \leq k_1 \leq k_2 \leq n$. Then $\Phi(c_1) \leq_{\mathcal{B}} \Phi(c_2)$ if and only if $\Phi(c_1) \cap \Phi(c_2) \neq \emptyset$ if and only if there exists a maximal chain $\alpha' \subset C$ which contains both c_1 and c_2 , if and only if $c_1 \leq c_2$ (since we already know that $\deg(c_1) \leq \deg(c_2)$). \Box

We deduce the proof of Theorem 4.3 that each local poset L_x in the Deligne complex of type \tilde{A}_{n-1} is a lattice.

Proof of Theorem 4.3 Let x denote a vertex of the Deligne complex X of type \tilde{A}_{n-1} . Without loss of generality, we may assume that x corresponds to the maximal proper parabolic subgroup $A(A_{n-1}) = \langle \sigma_1, \ldots, \sigma_{n-1} \rangle$, which is isomorphic to the Artin group of type A_{n-1} , ie the *n*-strand braid group. Since $A(A_{n-1})$ is a maximal parabolic subgroup of spherical type of the Euclidean Artin group $A(\tilde{A}_{n-1})$, the star of x in X identifies with the Deligne complex of the Artin group $A(A_{n-1})$. In particular, vertices in X adjacent to x may be identified with cosets of proper maximal parabolic subgroups of the braid group $A(A_{n-1})$. Furthermore, given two such cosets gP_i and hP_j for $g, h \in A(A_{n-1})$ and $1 \le i, j \le n-1$,

the corresponding vertices of X are adjacent if and only if $gP_i \cap hP_j \neq \emptyset$. As a consequence, the poset $L_{x,0}$ is isomorphic to the poset $\mathcal{B} \setminus \{0, 1\}$, and L_x is isomorphic to the poset \mathcal{B} . According to Theorem 5.10, we deduce that L_x is a lattice.

6 The thickening of a semilattice

We will now consider a generalization of Theorem 3.10 to the case of a semilattice. This will be useful to consider Euclidean buildings, Deligne complexes in type \tilde{C}_n and Artin groups of type FC. We start by recalling the definition of a flag poset.

Definition 6.1 A poset L is called *flag* if any three elements which are pairwise upper bounded have an upper bound.

Lemma 6.2 Let *L* denote a graded flag meet-semilattice with bounded rank. Then any family of elements of *L* which are pairwise upper bounded have a join.

Proof We first prove that every finite family of k elements $\{a_i\}_{1 \le i \le k}$ of L which are pairwise upper bounded have a join, by induction on the number of elements. By assumption, the property is true for k = 3. Fix $k \ge 4$, assume that the property is true for k-1, and consider k elements $\{a_i\}_{1 \le i \le k}$ of L which are pairwise upper bounded. Let b denote the join of a_{k-1} and a_k . The family $\{a_1, a_2, \ldots, a_{k-2}, b\}$ is pairwise upper bounded, so by assumption it has a join $c \in L$. Then c is the join of $\{a_i\}_{1 \le i \le k}$, which proves the induction.

Now consider an arbitrary family A of elements of L which are pairwise upper bounded. Since L has bounded rank, we may consider a finite subset $F \subset A$ such that the rank of the join b of F is maximal among all joins of finite subfamilies of A. For any $a \in A$, the join b_a of $F \cup \{a\}$ satisfies $b \leq b_a$, and the rank of b_a is at most the rank of b; hence $b_a = b$. We deduce that b is the join of A.

Theorem 6.3 Let *L* denote a graded poset with minimum 0 and with bounded rank such that:

- *L* is a meet-semilattice.
- L is flag.

Then the ℓ^{∞} orthoscheme realization of |L| is injective.

Proof Let us consider $\overline{1} = L \cup \{1\}$: it is a bounded graded lattice. Let us denote by $\overline{1} = |\overline{1}|$ the geometric realization of $\overline{1}$, and by $M = |L| \subset \overline{1}$ the geometric realization of L. Note that we consider $\overline{1}$ as a subset of $\overline{1}$, its vertex set. The orthoscheme realization of M is endowed with the induced length metric as a subspace of the ℓ^{∞} orthoscheme realization of $\overline{1}$.

If we consider the affine version $\bar{1}_{\mathbb{R}}$ of $\bar{1}$ over \mathbb{R} , then the elements $0_M = [(0, ..., 0), c] \in \bar{1}_{\mathbb{R}}$ and $1_M = [(1, ..., 1), c] \in \bar{1}_{\mathbb{R}}$ are such that $\bar{1}$ naturally identifies with the interval $I_{\bar{1}_{\mathbb{R}}}(0_M, 1_M)$ in $\bar{1}_{\mathbb{R}}$, as in the proof of Theorem 3.10. In particular, M and $\bar{1}$ are also posets.

4048

According to Theorem 3.8, the space $\bar{1}_{\mathbb{R}}$, with its standard ℓ^{∞} metric \bar{d} , is injective. According to Theorem 3.10, its interval $\bar{1} = I_{\bar{1}_{\mathbb{R}}}(0_M, 1_M)$ is injective. Let us denote by d the length metric of M.

Fix $\varepsilon > 0$, and consider the graph Γ_{ε} with vertex set M, and with an edge between $x, y \in M$ if $\overline{d}(x, y) \leq \varepsilon$. We will prove that Γ_{ε} is Helly, by proving that it is clique-Helly and that its triangle complex is simply connected.

Let $\sigma \subset \Gamma_{\varepsilon}$ denote a maximal clique: we claim that there exist $a, b \in \overline{1}$ such that σ is the intersection of an interval $I_{\overline{1}}(a, b)$ in $\overline{1}$ with M. Since σ is a subset of $\overline{1}_{\mathbb{R}}$ of diameter at most ε , there exist $a_0, b_0 \in \overline{1}_{\mathbb{R}}$ such that $\sigma \subset I_{\overline{1}_{\mathbb{R}}}(a_0, b_0)$ and $\overline{d}(a_0, b_0) \leq \varepsilon$. Since $\overline{1}$ is an interval, we deduce that there exist $a, b \in \overline{1} \cap I_{\overline{1}_{\mathbb{R}}}(a_0, b_0)$ such that $\sigma \subset I_{\overline{1}_{\mathbb{R}}}(a, b) = I_{\overline{1}}(a, b)$. In particular, we also have $\overline{d}(a, b) \leq \varepsilon$. Since σ is a maximal clique, we deduce that $\sigma = I_{\overline{1}}(a, b) \cap M$.

We will prove that Γ_{ε} is clique-Helly. Consider a family of pairwise intersecting maximal cliques $(\sigma_i)_{i \in I}$ in Γ_{ε} . For each $i \in I$, according to the previous paragraph, there exists $a_i \leq b_i$ in $\overline{1}$ such that $\sigma_i = I_{\overline{1}}(a_i, b_i) \cap M$. For each $i \in I$, let $m_i \in L$ minimal such that $a_i \leq m_i$. For each $i \neq j$ in I, since σ_i and σ_j intersect, the intersection $I_{\overline{1}}(a_i, b_i) \cap I_{\overline{1}}(a_j, b_j) \cap M$ is nonempty. In particular, the elements m_i and m_j have a common upper bound in L.

The family $(m_i)_{i \in I}$ is pairwise upper bounded. Since *L* is a graded flag semilattice with bounded rank, according to Lemma 6.2, this family has a join $m = \bigvee_{i \in I} m_i$ in *L*. In particular, the family $(\sigma_i \cap I_{\bar{1}}(0,m))_{i \in I}$ of intervals in the lattice $I_{\bar{1}}(0,m) = I_M(0,m)$ has a nonempty intersection. So the graph Γ_{ε} is clique-Helly.

We will now prove that the triangle complex of Γ_{ε} is simply connected. For each $t \in [0, 1]$, consider the map $\pi_t \colon M \to M$ sending each $x \in I_{\overline{1}}(0, m)$ to the point on the affine segment joining 0 to x at distance td(0, x) from 0. Note that π_t is 1–Lipschitz with respect to the distance d. Furthermore, if $t, t' \in [0, 1]$ are such that $|t - t'| \leq \varepsilon$, then for each $x \in M$ we have $d(\pi_t(x), \pi_{t'}(x)) \leq \varepsilon$. As a consequence, any combinatorial loop γ in Γ_{ε} may be homotoped in the triangle complex of Γ_{ε} to the loop $\pi_0(\gamma)$, which is the constant loop at 0. So we have proved that the triangle complex of Γ_{ε} is simply connected.

According to Theorem 1.12, we deduce that the graph Γ_{ε} is Helly. In particular, for each $\varepsilon > 0$, the metric space M is ε -coarsely injective. Since M is complete according to Theorem 1.11, we deduce by Lemma 1.13 that M is injective.

We will now give natural examples of such semilattices, which will be used in the sequel for Euclidean buildings and Deligne complexes of Euclidean type different from \tilde{A}_n .

Proposition 6.4 Let L_0 denote the vertex set of a (possibly nonthick) spherical building of type B_n for some $n \ge 1$, and let $L = L_0 \cup \{0\}$. In other words, L is the poset of subspaces of a polar space of projective dimension n - 1. Then L is a graded semilattice of rank n with minimum 0 such that any pairwise upper bounded subset of L has a join.

 s_1 s_2 s_3 s_{n-2} s_{n-1} s_n • ---- • · · · · • ---- • 4

Figure 4: The Dynkin diagram of type B_n .

Proof The semilattice property is obvious: if S, S' are subspaces of a polar space X, then $S \wedge S' = S \cap S'$. If $A \subset L$ are pairwise upper bounded in L, let $A_X = \bigcup_{S \in A} S \subset X$. For any $x, y \in A_X$, there exists a subspace of X containing x and y. According to [Tits 1974, 7.2.1], the subset A_X is contained in a subspace S_0 of X. Hence, for any $S \in A$, we have $S \subset S_0$. So the intersection of all subspaces of Xcontaining $\bigcup_{S \in A} S$ is the join of A.

Another application concerns the Artin group $A(B_n)$ of spherical type B_n , with the following Dynkin diagram (see Figure 4).

Let s_1, \ldots, s_n denote the standard generators of $A(B_n)$, with $s_{n-1}s_ns_{n-1}s_n = s_ns_{n-1}s_ns_{n-1}$. For each $1 \le i \le n$, let P_i denote the maximal proper standard parabolic subgroup of $A(B_n)$

$$P_i = \langle s_1, \ldots, s_{i-1} \rangle \times \langle s_{i+1}, \ldots, s_n \rangle$$

Let $L_0 = \{gP_i \mid g \in A(B_n), 1 \le i \le n\}$ and $L = L_0 \cup \{0\}$. We define an order relation on L as follows. First, define $0 \le gP_i$ for each $gP_i \in L$. Otherwise, for $g_1, g_2 \in A(B_n)$ and $1 \le k_1, k_2 \le n$, define $g_1P_{k_1} \le g_2P_{k_2}$ if $k_1 \le k_2$ and $g_1P_{k_1} \cap g_2P_{k_2} \ne \emptyset$.

Lemma 6.5 The relation \leq is a partial order on *L*.

Proof First note that \leq is antisymmetric: if $g_1 P_{k_1}, g_2 P_{k_2} \in L_0$ are such that $g_1 P_{k_1} \leq g_2 P_{k_2}$ and $g_2 P_{k_2} \leq g_1 P_{k_1}$, then $k_1 = k_2$, and $g_1 P_{k_1} \cap g_2 P_{k_1} \neq \emptyset$ implies that $g_1 P_{k_1} = g_2 P_{k_1}$.

Now we show that \leq is transitive: assume that $g_1 P_{k_1}, g_2 P_{k_2}, g_3 P_{k_3} \in L_0$ are such that $g_1 P_{k_1} \leq g_2 P_{k_2}$ and $g_2 P_{k_2} \leq g_3 P_{k_3}$. We deduce that $k_1 \leq k_2 \leq k_3, g_1 \in g_2 P_{k_2} P_{k_1}$ and $g_2 \in g_3 P_{k_3} P_{k_2}$, so $g_1 \in g_3 P_{k_3} P_{k_2} P_{k_1}$. Note that $\langle s_1, \dots, s_{k_2-1} \rangle \subset P_{k_3}$ and $\langle s_{k_2+1}, \dots, s_n \rangle \subset P_{k_1}$, so $P_{k_3} P_{k_2} P_{k_1} = P_{k_3} P_{k_1}$. In particular $g_1 \in g_3 P_{k_3} P_{k_1}$ and $k_1 \leq k_3$, so $g_1 P_{k_1} \leq g_3 P_{k_3}$.

Note that we are grateful to Luis Paris for his help in the following proof, notably the use of normal forms.

Proposition 6.6 L is a graded semilattice of rank n with minimum 0 such that any pairwise upper bounded subset of L has a join.

Proof Let t_1, \ldots, t_{2n-1} be the standard generators of the braid group $A(A_{2n-1})$. Consider the morphism

$$\phi: A(B_n) \to A(A_{2n-1}),$$

for all $1 \le i \le n-1$, $s_i \mapsto t_i t_{2n-i}$, $s_n \mapsto t_n$

This morphism ϕ is injective; see for instance [Dehornoy and Paris 1999; Michel 1999; Crisp 2000]. Let σ denote the involution of $A(A_{2n-1})$ defined for all $1 \le i \le 2n-1$ by $\sigma(t_i) = t_{2n-i}$. According to [Crisp 2000, Theorem 4], we know furthermore that the image of ϕ coincides with the fixed-point set of σ .

For each $1 \le i \le 2n - 1$, consider the standard proper maximal parabolic subgroup

$$Q_i = \langle t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_{2n-1} \rangle$$

of $A(A_{2n-1})$. Let M_0 denote the poset of cosets of maximal proper parabolic subgroups of $A(A_{2n-1})$, and $M = M_0 \cup \{0, 1\}$. We will define a poset map $\psi : L_0 \to M_0$.

For each $g \in A(B_n)$ and each $1 \le i \le n$, let us define $\psi(gP_i) = \phi(g)Q_i$. Since $\phi(P_i) \subset Q_i$, the map $\psi: L_0 \to M_0$ is well-defined. Assume that $gP_i \le hP_j$, ie $i \le j$ and $h^{-1}g \in P_jP_i$. Then $\phi(h)^{-1}\phi(g) \in \phi(P_j)\phi(P_i) \subset Q_jQ_i$, so $\phi(g)Q_i \le \phi(h)Q_j$. As a consequence, ψ is a rank-preserving injective poset map.

Note that the involution σ extends naturally to an order-reversing involution on M by letting $\sigma(gQ_i) = \sigma(g)Q_{2n-i}$. And for each $g \in A(B_n)$ and each $1 \le i \le n$, we have $\sigma(\psi(gP_i)) = \phi(g)Q_{2n-i}$.

We will now prove that *L* is a meet-semilattice. Fix $a, b \in A(B_n)$ and $1 \le i, j \le n$; we will prove that aP_i and bP_j have a meet in *L*. Consider the elements $\phi(a)Q_i$ and $\phi(b)Q_j$ in *M*: they have a meet γQ_k for some $\gamma \in A(A_{2n-1})$ and $0 \le k \le i, j$. Then as $\phi(a)Q_i \le \phi(a)Q_{2n-i}$ and $\phi(b)Q_j \le \phi(b)Q_{2n-j}$, we deduce that $\gamma Q_k \le \sigma(\gamma)Q_{2n-k}$.

So, up to the choice of $\gamma \in \gamma Q_k$, we may assume that $\sigma(\gamma) \in \gamma Q_{2n-k}$: let $q \in Q_{2n-k}$ such that $\sigma(\gamma) = \gamma q$. Since σ is an involution, we have $\gamma = \gamma q \sigma(q)$, so $q = \sigma(q)^{-1} \in Q_k \cap Q_{2n-k}$.

We claim that if $g \in A(A_{2n-1})$ is such that $\sigma(g) = g^{-1}$, there exists $h \in A(A_{2n-1})^+$ such that $g = h\sigma(h)^{-1}$. According to [Charney 1995], there exist unique $h, h' \in A(A_{2n-1})^+$ such that $g = hh'^{-1}$ and the right greatest common divisor of h and h' in the Garside monoid $A(A_{2n-1})^+$ is 1. Note that σ preserves the monoid $A(A_{2n-1})^+$; hence we have $g^{-1} = h'h^{-1}$ on one side, and $g^{-1} = \sigma(g) = \sigma(h)\sigma(h')^{-1}$ on the other side. By the uniqueness of h, h', we deduce that $h' = \sigma(h)$. Hence $g = h\sigma(h)^{-1}$.

Assume furthermore that $g \in Q_k \cap Q_{2n-k} = \langle t_1, \dots, t_{k-1} \rangle \times \langle t_k + 1, \dots, t_{2n-k-1} \rangle \times \langle t_{2n-k+1}, \dots, t_{2n-1} \rangle$. Then we can decompose g as a fraction $g = hh'^{-1}$ inside the parabolic subgroup $Q_k \cap Q_{2n-k}$: by the uniqueness of h, h', we deduce that $h, h' \in Q_k \cap Q_{2n-k}$.

According to the claim, there exists $q' \in Q_k \cap Q_{2n-k}$ such that $q = q'\sigma(q')^{-1}$. So, up to replacing γ with $\gamma' = \gamma q' \in \gamma Q_k$, we have

$$\sigma(\gamma') = \sigma(\gamma)\sigma(q') = \gamma q \sigma(q') = \gamma q' = \gamma'.$$

So we may assume that γ' is fixed by σ : according to [Crisp 2000, Theorem 4], we may consider $c \in A(B_n)$ such that $\phi(c) = \gamma'$. So we deduce that $cP_k \leq aP_i, bP_j$. Conversely, for any $c' \in A(B_n)$ and $k' \leq i, j$ such that $c'P_{k'} \leq aP_i, bP_j$, we have $\phi(c')Q_{k'} \leq \phi(a)Q_i \wedge \phi(b)Q_j = \phi(c)Q_k$, so $c'P_{k'} \leq cP_k$. We conclude that cP_k is the meet of aP_i and bP_j in *L*.

Algebraic & Geometric Topology, Volume 24 (2024)

We will now prove that any three pairwise upper bounded elements of *L* have an upper bound. Fix $g_1, g_2, g_3 \in A(B_n)$ and $1 \leq k_1, k_2, k_3 \leq n$ such that, for each $i \neq j$, the elements $g_i P_{k_i}$ and $g_j P_{k_j}$ have a join $g_{ij} P_{k_{ij}}$ in *L*. Consider the elements $\phi(g_{ij})Q_{k_{ij}}$ of *M*: they have a join γQ_k in *M*. Since, for each $1 \leq i, j \leq 3$, we have $\phi(g_i)Q_i \leq \phi(g_{ij})Q_{k_{ij}} \leq \phi(g_{ij})Q_{2n-k_{ij}} \leq \phi(g_j)Q_{2n-k_{ij}}$, we deduce that γQ_k is inferior to the meet $\sigma(\gamma)Q_{2n-k}$ of the elements $\phi(g_{ij})Q_{2n-k_{ij}}$. We deduce that $k \leq n$. Also, as in the previous paragraph, we deduce that we can choose $\gamma \in A(A_{2n-1})$ such that $\gamma = \phi(c)$ for some $c \in A(B_n)$. Hence cP_k is a common upper bound for the elements $(g_i P_{k_i})_{1 \leq i \leq 3}$.

Since L is graded with finite rank, we deduce that any family of pairwise upper bounded elements of L have a join. \Box

We believe that a similar statement holds for the Artin group of spherical type D_n , but we have no embedding into an Artin group of type A in order to use a similar proof.

7 Application to Euclidean buildings and the Deligne complex of other Euclidean types

In this section, we will prove that we can deduce from Theorem 4.4 an injective metric on Euclidean buildings and Deligne complexes of Euclidean types other than \tilde{A}_n .

More precisely, let us consider a simplicial complex X that is

- either a Euclidean building of type \tilde{B}_n , \tilde{C}_n or \tilde{D}_n ,
- or the Deligne complex of Euclidean type \tilde{C}_n .

Note that the Coxeter groups of types \tilde{B}_n and \tilde{D}_n may be considered as subgroups of the Coxeter group of Euclidean type \tilde{C}_n , spanned by reflections with respect to subarrangements of hyperplanes; see below for more details. As a consequence, if X is a Euclidean building of type \tilde{B}_n or \tilde{D}_n , we will consider it as a (possibly nonthick) Euclidean building of type \tilde{C}_n .

Let $W(\tilde{C}_n)$ denote the Euclidean Coxeter group of type \tilde{C}_n . Its Coxeter complex $\Sigma(\tilde{C}_n)$ identifies with \mathbb{R}^n , and reflections of $W(\tilde{C}_n)$ correspond to reflections with respect to hyperplanes

 $\{x_i = k \mid 1 \le i \le n, k \in \mathbb{Z}\}$ and $\{x_i \pm x_j = 2k \mid 1 \le i \ne j \le n, k \in \mathbb{Z}\}.$

Note that we can see the following subarrangement has type \tilde{B}_n :

 $\{x_i = 2k \mid 1 \le i \le n, k \in \mathbb{Z}\} \text{ and } \{x_i \pm x_j = 2k \mid 1 \le i \ne j \le n, k \in \mathbb{Z}\}.$

Also the following subarrangement has type \tilde{D}_n :

$$\{x_i \pm x_j = 2k \mid 1 \le i \ne j \le n, k \in \mathbb{Z}\}.$$

This justifies that, in the case of Euclidean buildings, we may assume that X is a (possibly nonthick) Euclidean building of type \tilde{C}_n .

The vertex set of $\Sigma(\tilde{C}_n)$ identifies with \mathbb{Z}^n , and a strict fundamental domain of the action of $W(\tilde{C}_n)$ on $\Sigma(\tilde{C}_n)$ is given by the standard orthoscheme simplex σ with vertices

$$v_0 = (0, \dots, 0), \quad v_1 = (1, 0, \dots, 0), \quad \dots, \quad v_n = (1, 1, \dots, 1).$$

Hence v_0, v_1, \ldots, v_n are representatives of the n+1 orbits of vertices of $\Sigma(\tilde{C}_n)$ under the action of $W(\tilde{C}_n)$. This enables us to define a type function τ on vertices of $\Sigma(\tilde{C}_n)$:

 $\tau: \Sigma(\tilde{C}_n)^{(0)} \to \{0, 1, \dots, n\}, \quad g \cdot v_i \mapsto i.$

More generally, we can define a partial order on vertices of $\Sigma(\tilde{C}_n)$: say that v < v' if v, v' are adjacent vertices and $\tau(v) < \tau(v')$. We can therefore also view $\Sigma(\tilde{C}_n)$ as the geometric realization of the poset of its vertices.

We will now see how to extend the type function and the partial order on vertices of X.

Proposition 7.1 Assume that X is a Euclidean building of type \tilde{C}_n or the Deligne complex of Euclidean type \tilde{C}_n . There exists a type function $\tau: X^{(0)} \to \{0, 1, ..., n\}$ such that adjacent vertices of X have different types. Moreover, let us define a partial order on vertices of X by setting v < v' if v and v' are adjacent in X and $\tau(v) < \tau(v')$. Then X is the geometric realization of the poset of its vertices.

Proof Let us first consider the case where X is a Euclidean building of type \tilde{C}_n . Given any vertex v of X, consider any apartment $A \subset X$ containing v, and let us define $\tau(v)$ as defined with respect to the apartment $A \simeq \Sigma(\tilde{C}_n)$. Since two apartments containing v differ by an element of the Weyl group $W(\tilde{C}_n)$, which preserves the type, we deduce that τ is well-defined on $X^{(0)}$. Similarly, given any two adjacent vertices v, v' in X, say that v < v' if $\tau(v) < \tau(v')$.

We will see that this relation is actually transitive on vertices of X: assume that three vertices v_1, v_2, v_3 of X satisfy $v_1 < v_2$ and $v_2 < v_3$. Then the link of v_2 is isomorphic to the join of two spherical buildings of types $B_{\tau(v_2)}$ and $B_{n-\tau(v_2)}$. Hence we see that v_1 is adjacent to v_3 in X, so $v_1 < v_3$.

Let us now consider the case where X is the Deligne complex of Euclidean type \tilde{C}_n . Let $s_0, s_1, \ldots, s_{n-1}, s_n$ denote the standard generators of the Artin group $A(\tilde{C}_n)$. For each $0 \le i \le n$, consider the maximal spherical-type standard parabolic subgroup

$$P_i = \langle s_0, s_1, \ldots, s_{i-1}, s_{i+1}, \ldots, s_n \rangle \simeq A(B_i) \times A(B_{n-i}).$$

For each vertex gP_i in X, where $g \in A(\tilde{C}_n)$ and $0 \le i \le n$, let us define $\tau(gP_i) = i$. Given any two vertices gP_i, hP_j of X, say that $gP_i < hP_j$ if they are adjacent in X (ie $gP_i \cap hP_j \ne \emptyset$) and i < j. As in Lemma 6.5, one checks that it is a partial order.

We will endow X with the piecewise ℓ^{∞} orthoscheme metric d given by the geometric realization of its poset of vertices. We will now describe the local structure of X at any vertex, starting with a general statement about orthoscheme complexes of posets.

Algebraic & Geometric Topology, Volume 24 (2024)

Lemma 7.2 Let *L* denote a graded poset, with an element $v \in L$ comparable to every element of *L*. Let $L^+ = \{w \in L \mid w \ge v\}$ and $L^- = \{w \in L \mid w \le v\}$. Then the ℓ^{∞} orthoscheme realization of *L* is locally isometric at *v* to the ℓ^{∞} product of the ℓ^{∞} orthoscheme realizations of L^+ and L^- .

Proof Since the ℓ^{∞} orthoscheme realization of *L* is obtained as a union of orthoschemes, it is sufficient to prove the result when *L* is a chain.

In other words, consider the standard ℓ^{∞} *n*-orthoscheme $C_n = \{x \in \mathbb{R}^n \mid 1 \le x_1 \le x_2 \ge \cdots \ge x_n \ge 0\}$, with vertices $v_i = (1, \dots, 1, 0, \dots, 0)$ with *i* ones and n - i zeros for $0 \le i \le n$. We fix a particular vertex $v = v_j$ for some $0 \le j \le n$, and we want to describe locally *C* around *v*. The orthoscheme C_n is locally isometric at *v* to the space

$$T = \{ x \in \mathbb{R}^n \mid 1 \ge x_1 \ge x_2 \ge \cdots \ge x_j \text{ and } x_{j+1} \ge x_{j+2} \ge \cdots \ge x_n \ge 0 \},\$$

which is isometric to the ℓ^{∞} product of $T^- = \{(x_1, \dots, x_j) \in \mathbb{R}^j \mid 1 \ge x_1 \ge x_2 \ge \dots \ge x_j\}$ and $T^+ = \{(x_{j+1}, \dots, x_n) \in \mathbb{R}^{n-j} \mid x_{j+1} \ge x_{j+2} \ge \dots \ge x_n \ge 0\}.$

The space T^+ is locally isometric at v to the vertex (1, 1, ..., 1) in the *j*-orthoscheme C_j , and T^- is locally isometric at v to the vertex (0, 0, ..., 0) in the (n-j)-orthoscheme C_{n-j} .

In conclusion, if *L* is a chain $v_0 < v_1 < \cdots < v = v_j < v_n$, then the geometric realization of |L| is locally isometric at *v* to the product of the geometric realizations of the chains $L^- = (v_0 < v_1 < \cdots < v_j = v)$ and $L^+ = (v = v_j < v_{j+1} < \cdots < v_n)$.

Proposition 7.3 Assume that X is a Euclidean building of type \tilde{C}_n or the Deligne complex of Euclidean type \tilde{C}_n . Fix a vertex $v \in X$ of type $\tau(v) = i \in \{0, 1, ..., n\}$.

There exist posets $L_{1,i}$ (resp. $L'_{0,n-i}$), which are either the poset of vertices of a spherical building of type B_i (resp. B_{n-i}) or the poset of maximal proper parabolic subgroups of the Artin group of spherical type B_i (resp. B_{n-i}). Let us define the posets $L_i = \{1\} \cup L_{1,i}$ and $L'_{n-i} = L'_{0,n-i} \cup \{0\}$, and let us consider the geometric realizations $|L_i|$ and $|L'_{n-i}|$, endowed with the piecewise ℓ^{∞} orthoscheme metrics.

Then X is locally isometric at v to the ℓ^{∞} direct product $|L_i| \times |L'_{n-i}|$.

Proof The space X is locally isometric at v to the ℓ^{∞} orthoscheme realization of the poset L_v of vertices of X comparable to v. The poset L_v is the disjoint union of $L_{1,i} = \{w \in L_v \mid w < v\} \sqcup \{v\} \sqcup L'_{0,n-i} = \{w \in L_v \mid w > v\}$, such that $L_{1,i} < \{v\} < L'_{0,n-i}$. We may identify the poset $L_i = L_{1,i} \cup \{1\}$ with the interval $L_{1,i} \cup \{v\}$ in L_v . Similarly, we may identify the poset $L'_{n-i} = L'_{0,n-i} \cup \{0\}$ with the interval $L'_{0,n-i} \cup \{v\}$ in L_v . According to Lemma 7.2, the orthoscheme realization $|L_v|$ of L_v is locally isometric at v to the direct ℓ^{∞} product $|L_i| \times |L'_{n-i}|$ of the orthoscheme realizations of L_i and L'_{n-i} .

In case X is a Euclidean building to type \tilde{C}_n , note that $L_{1,i}$ is the vertex set of a spherical building of type B_i (with reversed order), and that $L'_{0,n-i}$ is the vertex set of a spherical building of type B_{n-i} .

4054

In the case X is the Deligne complex of type \tilde{C}_n , note that $L_{1,i}$ is the poset of proper parabolic subgroups of the Artin group of type B_i (with reversed order), and that $L'_{0,n-i}$ is the poset of proper parabolic subgroups of the Artin group of type B_{n-i} .

We can now prove the following.

Theorem 7.4 Any Euclidean building of type \tilde{B}_n , \tilde{C}_n or \tilde{D}_n , or the Deligne complex of type \tilde{C}_n , endowed with the piecewise ℓ^{∞} metric *d*, is injective.

Proof According to Proposition 7.3, the space X is locally isometric to a product of orthoscheme complexes of the type |L|, where $L = L_0 \cup \{0\}$ is a poset, and L_0 is either the poset of vertices of a spherical building of type B_i or the poset of maximal proper parabolic subgroups of the Artin group of spherical type B_i for some $0 \le i \le n$.

According to Propositions 6.4 and 6.6, we know that in each case L is a graded meet-semilattice with minimum 0 of rank *i* such that any upper bounded subset has a join. According to Theorem 6.3, we deduce that |L| is injective.

We know X is uniformly locally injective. According to Theorem 1.14, we deduce that X is injective. \Box

We can also deduce that the thickening is a Helly graph.

Corollary 7.5 The thickening of the vertex set of any Euclidean building of type \tilde{B}_n , \tilde{C}_n or \tilde{D}_n , or the extended Deligne complex of type \tilde{C}_n , is a Helly graph.

Proof Let X denote either a Euclidean building of type \tilde{B}_n , \tilde{C}_n or \tilde{D}_n , or the extended Deligne complex of type \tilde{C}_n . We will see that X satisfies the assumptions of Theorem 1.15.

Since simplices of X have a well-defined order, we see that X is a simplicial complex with ordered simplices. For each maximal simplex σ of X, the minimal and maximal vertices of σ form a maximal edge in X. Therefore, X has maximal edges as in Definition 1.9. According to Theorem 1.15, we deduce that the thickening of X is a Helly graph.

One can also deduce another proof of the result of Huang and Osajda that FC-type Artin groups are Helly (see [Huang and Osajda 2021, Theorem 5.8]), with moreover explicit Helly and injective models.

Theorem 7.6 Let $A = A(\Gamma)$ denote an FC-type Artin group, with standard generating set *S*, and let \leq_L denote the standard prefix order on *A* with respect to *S*.

• Consider the graph *Y*, with vertex set *A*, with an edge between $g, h \in A$ if there exists $a \in A$ and a spherical subset $T \subset S$ such that $a \leq_L g, h \leq_L a \Delta_T$, where Δ_T denotes the standard Garside element of A(T). Then *Y* is a Helly graph.

• Consider the simplicial complex X, with vertex set A, with a k-simplex for each chain $g_0 <_L g_1 <_L \cdots <_L g_k \leq_L g_0 \Delta_T$ for some spherical subset $T \subset S$. Endow X with the standard ℓ^{∞} orthoscheme metric. Then X is an injective metric space.

Proof Let us start by proving that X is injective. Note that X can be described as the union of (possibly not connected) complexes X_T , for $T \subset S$ spherical, by restricting the spherical subsets to be contained in T. So X is locally isometric to the geometric realization of the poset $L = \bigcup_{T \subset S} \text{ spherical}[1, \Delta_T]$. This poset is graded, with minimum, with bounded rank, and is a meet-semilattice.

Moreover, the flag condition from the FC-type Artin group A translates into the flag condition for the poset L. Indeed, consider x_1, x_2, x_3 which are pairwise upper bounded. Let $T_1, T_2, T_3 \subset S$ denote the supports of x_1, x_2, x_3 respectively. By assumption, for each $i \neq j$, the subset $T_i \cup T_j$ is spherical. As a consequence, the subset $T = T_1 \cup T_2 \cup T_3$ is a complete subset of S. According to the FC-type condition, we deduce that T is spherical. Hence $x_1, x_2, x_3 \leq_L \Delta_T$. So L is a flag poset.

According to Theorem 6.3, we deduce that |L| is injective. So X is uniformly locally injective and according to Theorem 1.14, we deduce that X is injective.

We now turn to the proof that Y is a Helly graph, as in the proof of Corollary 7.5. Since simplices of X have a well-defined order, we see that X is a simplicial complex with ordered simplices. For each maximal simplex σ of X, the minimal and maximal vertices of σ form a maximal edge in X. Therefore X has maximal edges. According to Theorem 1.15, we deduce that the thickening Y of X is a Helly graph. \Box

8 Bicombings on Deligne complexes in types $\tilde{A_n}$ and \tilde{C}_n

We now see that, in the Deligne complex of spherical types A_n , B_n and Euclidean types \tilde{A}_n , \tilde{C}_n , we may find a convex bicombing.

Theorem 8.1 Let X denote the Deligne complex of the Artin group A of spherical type A_n , B_n or Euclidean type \tilde{A}_n , \tilde{C}_n . There is a metric d_X on X that admits a convex, consistent, reversible geodesic bicombing σ . Moreover, d_X and σ are invariant under A.

Proof The statement for types A_n and \tilde{A}_n is Theorem 4.6.

Assume that X is the Deligne complex of type B_n . We have seen in the proof of Proposition 6.6 that X may be realized as the fixed-point subspace of the Deligne complex Y of type A_{2n-1} for an involution s. Note that, since s is order-reversing, s induces an isometry of Y. According to Theorem 4.6, there exists a unique reversible, convex, consistent, geodesic bicombing σ_Y on Y for a metric d_Y . Since σ_Y is unique, we deduce that s preserves σ_Y , and that the fixed-point subspace X is σ -stable. Let d_X denote the restriction of d_Y to X, and let σ_X denote the restriction of σ_Y to X. We deduce that σ_X is a convex, consistent, reversible geodesic bicombing on (X, d_X) . Moreover, d_X and σ_X are invariant under A.

Assume that X is the Deligne complex of type \tilde{C}_n . According to [Digne 2012, Theorem 5.2] (it is also a consequence of Corollary 8.5), we see that X may be realized as the fixed-point subspace of the Deligne complex Y of type \tilde{A}_{2n-1} for an involution s. Following the same arguments as in the type B_n case, we conclude that there exists a metric d_X on X and a convex, consistent, reversible geodesic bicombing σ_X on (X, d_X) that are both invariant under A.

We will describe several consequences we can derive from the fact that the Deligne complex has a consistent convex bicombing, which were usually known when the Deligne complex had a CAT(0) metric.

Corollary 8.2 (Okonek) The Deligne complex X of Euclidean type \tilde{A}_n or \tilde{C}_n is contractible. In particular, the $K(\pi, 1)$ conjecture holds in these cases.

Proof Metric spaces with a convex bicombing are contractible.

The proof of Morris-Wright [2021] for the intersection of parabolic subgroups in FC-type Artin groups adapts directly to our situation. It relies mainly on the result by Cumplido et al. [2019] that in a spherical-type Artin group, the intersection of parabolic subgroups is a parabolic subgroup.

Corollary 8.3 Let A denote the Artin group of Euclidean type \tilde{A}_n or \tilde{C}_n . The intersection of any family of parabolic subgroups of A is a parabolic subgroup.

Proof Note that any proper parabolic subgroup of an Artin group of Euclidean type has spherical type. Since the proof of [Morris-Wright 2021, Theorem 3.1] uses only the existence of an A-equivariant consistent geodesic bicombing on X, it adapts to these cases.

The results by Godelle [2007] describing centralizers and normalizers of parabolic subgroups also adapt to our case.

Corollary 8.4 Let A denote the Artin group of Euclidean type \tilde{A}_n or \tilde{C}_n . Then A satisfies properties (\star), ($\star\star$) and ($\star\star\star$) from [Godelle 2007]; notably, for any subset $X \subset S$, we have

 $\operatorname{Com}_{A}(A_{X}) = N_{A}(A_{X}) = A_{X} \cdot QZ_{A}(X),$

where the quasicentralizer of X is $QZ_A(X) = \{g \in A \mid g \cdot X = X\}.$

Note that Godelle's property $(\star\star)$ stating that a parabolic subgroup *P* is contained in a parabolic subgroup *Q*, then *P* is a parabolic subgroup of *Q* has been proved by Blufstein and Paris [2023].

Proof The proof of [Godelle 2007, Theorem 3.1] uses only the following properties of the Deligne complex, which hold for the Deligne complex X:

- There exists an A-equivariant consistent geodesic bicombing σ_X on X.
- Closed cells of X are σ_X -stable.

These assumptions are satisfied.

Note that Cumplido uses these properties of parabolic subgroups to solve the conjugacy stability problem for parabolic subgroups; see [Cumplido 2022, Theorem 14].

4056

The results by Crisp [2000] about symmetrical subgroups of Artin groups also extend to our case.

Corollary 8.5 Let A denote the Artin group of Euclidean type \tilde{A}_n or \tilde{C}_n . For any group G of symmetries of the Artin system, the fixed-point subgroup A^G is isomorphic to an Artin group.

For the explicit description of the Artin group A^G , we refer the reader to [Crisp 2000].

Proof The proof of [Crisp 2000, Theorem 23] uses only the following properties of the Deligne complex X:

- There exists an A-equivariant consistent geodesic bicombing σ_X on X.
- Subsets of X which are σ_X -stable are contractible.

Let us check the last condition: if $C \subset X$ is a σ_X -stable subset, then σ_X restricts to a convex bicombing on C, which implies that C is contractible.

One particular case of interest is when A is of Euclidean type \tilde{A}_{2n-1} for some $n \ge 1$, and the group G of symmetries of the Artin system (the 2n-cycle) is generated by a symmetry of the cycle defining A. We recover the result from [Digne 2012] that the fixed-point subgroup is isomorphic to the Artin group of type \tilde{C}_n .

For any divisor 1 < k < n of *n*, one may also consider the group *G* of symmetries of the Artin system of type \tilde{A}_{n-1} (the *n*-cycle) generated by a rotation of *k*. The fixed-point subgroup is then isomorphic to the Artin group of Euclidean type \tilde{A}_{k-1} . When we see $A(\tilde{A}_{n-1})$ as the group of braids with *n* strands on the annulus, we may think of the fixed-point subgroup as the subgroup of braids which are invariant by a rotation of the annulus of angle $\frac{2\pi k}{n}$.

References

- [Alonso and Bridson 1995] JM Alonso, MR Bridson, Semihyperbolic groups, Proc. Lond. Math. Soc. 70 (1995) 56–114 MR Zbl
- [Basso 2024] G Basso, Extending and improving conical bicombings, Enseign. Math. 70 (2024) 165–196 MR Zbl
- [Bessis 2006] D Bessis, A dual braid monoid for the free group, J. Algebra 302 (2006) 55–69 MR Zbl
- [Birman et al. 1998] J Birman, K H Ko, S J Lee, A new approach to the word and conjugacy problems in the braid groups, Adv. Math. 139 (1998) 322–353 MR Zbl
- [Blufstein and Paris 2023] MA Blufstein, L Paris, Parabolic subgroups inside parabolic subgroups of Artin groups, Proc. Amer. Math. Soc. 151 (2023) 1519–1526 MR Zbl
- [Bourbaki 2002] N Bourbaki, Lie groups and Lie algebras, chapters IV-VI, Springer (2002) MR Zbl
- [Bowditch 2020] **BH Bowditch**, *Median and injective metric spaces*, Math. Proc. Cambridge Philos. Soc. 168 (2020) 43–55 MR Zbl

- [Brady and McCammond 2010] T Brady, J McCammond, Braids, posets and orthoschemes, Algebr. Geom. Topol. 10 (2010) 2277–2314 MR Zbl
- [Bridson and Haefliger 1999] **M R Bridson**, **A Haefliger**, *Metric spaces of non-positive curvature*, Grundl. Math. Wissen. 319, Springer (1999) MR Zbl
- [Bruhat and Tits 1972] **F Bruhat**, **J Tits**, *Groupes réductifs sur un corps local*, Inst. Hautes Études Sci. Publ. Math. 41 (1972) 5–251 MR Zbl
- [Chalopin et al. 2020a] J Chalopin, V Chepoi, A Genevois, H Hirai, D Osajda, *Helly groups*, preprint (2020) arXiv 2002.06895
- [Chalopin et al. 2020b] J Chalopin, V Chepoi, H Hirai, D Osajda, *Weakly modular graphs and nonpositive curvature*, Mem. Amer. Math. Soc. 1309, Amer. Math. Soc., Providence, RI (2020) MR
- [Charney 1995] **R Charney**, *Geodesic automation and growth functions for Artin groups of finite type*, Math. Ann. 301 (1995) 307–324 MR Zbl
- [Charney 2004] **R** Charney, *The Deligne complex for the four-strand braid group*, Trans. Amer. Math. Soc. 356 (2004) 3881–3897 MR Zbl
- [Charney 2008] **R** Charney, *Problems related to Artin groups*, preprint, Amer. Inst. Math. (2008) Available at https://people.brandeis.edu/~charney/papers/Artin_probs.pdf
- [Charney and Davis 1995a] **R** Charney, MW Davis, *Finite* $K(\pi, 1)s$ for Artin groups, from "Prospects in topology", Ann. of Math. Stud. 138, Princeton Univ. Press (1995) 110–124 MR Zbl
- [Charney and Davis 1995b] **R** Charney, **M** W Davis, *The* $K(\pi, 1)$ –*problem for hyperplane complements associated to infinite reflection groups*, J. Amer. Math. Soc. 8 (1995) 597–627 MR Zbl
- [Crisp 2000] J Crisp, Symmetrical subgroups of Artin groups, Adv. Math. 152 (2000) 159–177 MR Zbl
- [Cumplido 2022] **M Cumplido**, *The conjugacy stability problem for parabolic subgroups in Artin groups*, Mediterr. J. Math. 19 (2022) art. id. 237 MR Zbl
- [Cumplido et al. 2019] M Cumplido, V Gebhardt, J González-Meneses, B Wiest, On parabolic subgroups of Artin–Tits groups of spherical type, Adv. Math. 352 (2019) 572–610 MR Zbl
- [Dehornoy and Paris 1999] **P Dehornoy**, **L Paris**, *Gaussian groups and Garside groups, two generalisations of Artin groups*, Proc. Lond. Math. Soc. 79 (1999) 569–604 MR Zbl
- [Dehornoy et al. 2015] **P Dehornoy, F Digne, E Godelle, D Krammer, J Michel**, *Foundations of Garside theory*, EMS Tracts in Math. 22, Eur. Math. Soc., Zürich (2015) MR Zbl
- [Deligne 1972] **P Deligne**, *Les immeubles des groupes de tresses généralisés*, Invent. Math. 17 (1972) 273–302 MR Zbl
- [Descombes and Lang 2015] **D Descombes**, **U Lang**, *Convex geodesic bicombings and hyperbolicity*, Geom. Dedicata 177 (2015) 367–384 MR Zbl
- [Descombes and Lang 2016] **D Descombes**, **U Lang**, *Flats in spaces with convex geodesic bicombings*, Anal. Geom. Metr. Spaces 4 (2016) 68–84 MR Zbl
- [Digne 2012] **F Digne**, A Garside presentation for Artin–Tits groups of type \tilde{C}_n , Ann. Inst. Fourier (Grenoble) 62 (2012) 641–666 MR Zbl
- [Dougherty et al. 2020] **M Dougherty**, **J McCammond**, **S Witzel**, *Boundary braids*, Algebr. Geom. Topol. 20 (2020) 3505–3560 MR Zbl

- [Dress 1984] **A W M Dress**, *Trees, tight extensions of metric spaces, and the cohomological dimension of certain groups: a note on combinatorial properties of metric spaces*, Adv. Math. 53 (1984) 321–402 MR Zbl
- [Godelle 2007] E Godelle, Artin–Tits groups with CAT(0) Deligne complex, J. Pure Appl. Algebra 208 (2007) 39–52 MR Zbl
- [Godelle and Paris 2012] E Godelle, L Paris, *Basic questions on Artin–Tits groups*, from "Configuration spaces", CRM Series 14, Ed. Norm., Pisa (2012) 299–311 MR Zbl
- [Haettel 2022a] **T Haettel**, *Injective metrics on buildings and symmetric spaces*, Bull. Lond. Math. Soc. 54 (2022) 2297–2313 MR Zbl
- [Haettel 2022b] **T Haettel**, *A link condition for simplicial complexes, and CUB spaces*, preprint (2022) arXiv 2211.07857
- [Haettel and Huang 2024] **T Haettel**, **J Huang**, *Lattices*, *Garside structures and weakly modular graphs*, J. Algebra 656 (2024) 226–258 MR Zbl
- [Haettel and Osajda 2021] **T Haettel**, **D Osajda**, *Locally elliptic actions, torsion groups, and nonpositively curved spaces*, preprint (2021) arXiv 2110.12431
- [Haettel et al. 2016] **T Haettel**, **D Kielak**, **P Schwer**, *The 6–strand braid group is* CAT(0), Geom. Dedicata 182 (2016) 263–286 MR
- [Haettel et al. 2023] T Haettel, N Hoda, H Petyt, Coarse injectivity, hierarchical hyperbolicity and semihyperbolicity, Geom. Topol. 27 (2023) 1587–1633 MR Zbl
- [Hoda 2023] N Hoda, Crystallographic Helly groups, Bull. Lond. Math. Soc. 55 (2023) 2991–3011 MR Zbl
- [Huang and Osajda 2021] J Huang, D Osajda, *Helly meets Garside and Artin*, Invent. Math. 225 (2021) 395–426 MR Zbl
- [Jeong 2023] S G Jeong, *The seven-strand braid group is* CAT(0), Manuscripta Math. 171 (2023) 563–581 MR Zbl
- [Kasprowski and Rüping 2017] **D Kasprowski**, **H Rüping**, *The Farrell–Jones conjecture for hyperbolic and* CAT(0)–*groups revisited*, J. Topol. Anal. 9 (2017) 551–569 MR
- [Lang 2013] U Lang, Injective hulls of certain discrete metric spaces and groups, J. Topol. Anal. 5 (2013) 297–331 MR Zbl
- [Leary and Minasyan 2021] IJ Leary, A Minasyan, Commensurating HNN extensions: nonpositive curvature and biautomaticity, Geom. Topol. 25 (2021) 1819–1860 MR
- [van der Lek 1983] **H van der Lek**, *The homotopy type of complex hyperplane complements*, PhD thesis, Radboud University (1983) Available at https://repository.ubn.ru.nl/handle/2066/148301
- [McCammond 2006] J McCammond, An introduction to Garside structures, preprint (2006) Available at https://web.math.ucsb.edu/~mccammon/current/artin/0intro/intro-garside.pdf
- [McCammond 2017] J McCammond, *The mysterious geometry of Artin groups*, Winter Braids Lect. Notes 4 (2017) art. id. 1 MR Zbl
- [McCammond and Sulway 2017] J McCammond, R Sulway, Artin groups of Euclidean type, Invent. Math. 210 (2017) 231–282 MR Zbl
- [Michel 1999] J Michel, A note on words in braid monoids, J. Algebra 215 (1999) 366–377 MR Zbl
- [Miesch 2014] **B Miesch**, *Injective metrics on cube complexes*, preprint (2014) arXiv 1411.7234

- [Miesch 2017] **B Miesch**, *The Cartan–Hadamard theorem for metric spaces with local geodesic bicombings*, Enseign. Math. 63 (2017) 233–247 MR Zbl
- [Möller et al. 2024] **P Möller, L Paris, O Varghese**, *On parabolic subgroups of Artin groups*, Israel J. Math. 261 (2024) 809–840 MR Zbl
- [Morris-Wright 2021] **R Morris-Wright**, *Parabolic subgroups in FC-type Artin groups*, J. Pure Appl. Algebra 225 (2021) art. id. 106468 MR Zbl
- [Nachbin 1950] L Nachbin, A theorem of the Hahn–Banach type for linear transformations, Trans. Amer. Math. Soc. 68 (1950) 28–46 MR Zbl
- [Okonek 1979] **C Okonek**, *Das K*(π , 1)–*Problem für die affinen Wurzelsysteme vom Typ A_n,C_n*, Math. Z. 168 (1979) 143–148 MR
- [Osajda and Valiunas 2024] **D** Osajda, M Valiunas, *Helly groups, coarsely Helly groups, and relative hyperbolicity*, Trans. Amer. Math. Soc. 377 (2024) 1505–1542 MR Zbl
- [Paolini and Salvetti 2021] **G Paolini**, **M Salvetti**, *Proof of the* $K(\pi, 1)$ *conjecture for affine Artin groups*, Invent. Math. 224 (2021) 487–572 MR Zbl
- [Świątkowski 2006] J Świątkowski, *Regular path systems and (bi)automatic groups*, Geom. Dedicata 118 (2006) 23–48 MR Zbl
- [Tits 1966] J Tits, Normalisateurs de tores, I: Groupes de Coxeter étendus, J. Algebra 4 (1966) 96-116 MR Zbl
- [Tits 1974] **J Tits**, *Buildings of spherical type and finite BN-pairs*, Lecture Notes in Math. 386, Springer (1974) MR Zbl

Institut Montpelliérain Alexander Grothendieck, Université de Montpellier, CNRS Montpellier, France

IRL 3457, CRM-CNRS, Université de Montréal Montréal, Canada

thomas.haettel@umontpellier.fr

Received: 15 February 2023 Revised: 12 September 2023



ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre	Kathryn Hess
etnyre@math.gatech.edu	kathryn.hess@epfl.ch
Georgia Institute of Technology	École Polytechnique Fédérale de Lausanne

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Steven Boyer	Université du Québec à Montréal cohf@math.rochester.edu	Robert Lipshitz	University of Oregon lipshitz@uoregon.edu
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Norihiko Minami	Yamato University minami.norihiko@yamato-u.ac.jp
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Robert Oliver	Université Paris 13 bobol@math.univ-paris13.fr
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futer	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Vesna Stojanoska	Univ. of Illinois at Urbana-Champaign vesna@illinois.edu
Ian Hambleton	McMaster University ian@math.mcmaster.ca	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Chris Wendl	Humboldt-Universität zu Berlin wendl@math.hu-berlin.de
Daniel Isaksen	Wayne State University isaksen@math.wayne.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Thomas Koberda	University of Virginia thomas.koberda@virginia.edu	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Markus Land	LMU München markus.land@math.lmu.de		

See inside back cover or msp.org/agt for submission instructions.

ł

The subscription price for 2024 is US \$705/year for the electronic version, and \$1040/year (+\$70, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by Mathematical Reviews, Zentralblatt MATH, Current Mathematical Publications and the Science Citation Index.

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, c/o Department of Mathematics, University of California, 798 Evans Hall #3840, Berkeley, CA 94720-3840.

AGT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY mathematical sciences publishers nonprofit scientific publishing https://msp.org/ © 2024 Mathematical Sciences Publishers

ALGEBRAIC

& **GEOMETRIC TOPOLOGY**

Volume 24 Issue 7 (pages 3571–4137) 2024 Geography of bilinearized Legendrian contact homology FRÉDÉRIC BOURGEOIS and DAMIEN GALANT The deformation spaces of geodesic triangulations of flat tori YANWEN LUO, TIANQI WU and XIAOPING ZHU Finite presentations of the mapping class groups of once-stabilized Heegaard splittings DAIKI IGUCHI On the structure of the top homology group of the Johnson kernel IGOR A SPIRIDONOV The Heisenberg double of involutory Hopf algebras and invariants of closed 3-manifolds SERBAN MATEI MIHALACHE, SAKIE SUZUKI and YUJI TERASHIMA A closed ball compactification of a maximal component via cores of trees GIUSEPPE MARTONE, CHARLES OUYANG and ANDREA TAMBURELLI An algorithmic discrete gradient field and the cohomology algebra of configuration spaces of two points on complete graphs EMILIO J GONZÁLEZ and JESÚS GONZÁLEZ Spectral diameter of Liouville domains PIERRE-ALEXANDRE MAILHOT Classifying rational G-spectra for profinite GDAVID BARNES and DANNY SUGRUE An explicit comparison between 2-complicial sets and Θ_2 -spaces JULIA E BERGNER, VIKTORIYA OZORNOVA and MARTINA ROVELLI On products of beta and gamma elements in the homotopy of the first Smith-Toda spectrum KATSUMI SHIMOMURA and MAO-NO-SUKE SHIMOMURA Phase transition for the existence of van Kampen 2-complexes in random groups TSUNG-HSUAN TSAI A qualitative description of the horoboundary of the Teichmüller metric AITOR AZEMAR Vector fields on noncompact manifolds TSUYOSHI KATO, DAISUKE KISHIMOTO and MITSUNOBU TSUTAYA Smallest nonabelian quotients of surface braid groups CINDY TAN Lattices, injective metrics and the $K(\pi, 1)$ conjecture THOMAS HAETTEL The real-oriented cohomology of infinite stunted projective spaces WILLIAM BALDERRAMA Fourier transforms and integer homology cobordism MIKE MILLER EISMEIER Profinite isomorphisms and fixed-point properties MARTIN R BRIDSON Slice genus bound in DTS^2 from *s*-invariant **QIUYU REN**

Relatively geometric actions of Kähler groups on CAT(0) cube complexes

COREY BREGMAN, DANIEL GROVES and KEJIA ZHU